



Regulation of Transcription






- activation of transcription factors (TF) - repressors and/or activators
- binding of TF(s) to DNA targets (~10 bp)
- enhance transcription by recruiting RNA polymerase (RNAP) to its target ("promoter")

- one TF can regulate (activate or repress) multiple promoters
- some promoters regulated by multiple TFs
- regulated gene may encode another regulator

PROKARYOTES






STATE	ENERGY	MULTIPLICITY	WEIGHT
<p>(A)</p> 	$L\varepsilon_{\text{sol}}$	$\frac{\Omega!}{L!(\Omega-L)!} \approx \frac{\Omega^L}{L!}$	$\frac{\Omega^L}{L!} e^{-\beta L\varepsilon_{\text{sol}}}$
<p>(B)</p> 	$(L-1)\varepsilon_{\text{sol}} + \varepsilon_{\text{b}}$	$\frac{\Omega!}{(L-1)!(\Omega-L+1)!} \approx \frac{\Omega^{L-1}}{(L-1)!}$	$\frac{\Omega^{L-1}}{(L-1)!} e^{-\beta[(L-1)\varepsilon_{\text{sol}} + \varepsilon_{\text{b}}]}$

Measuring regulatory functions quantitatively

the setup:




assumptions:

- fluorescence ~ concentration
- well isolated system


need to:

- calibrate expression
- quantify parameters
- make sure degradation is negligible
- measure cl and gene expression simultaneously


checks:



results:




Measuring regulatory functions quantitatively



Measuring regulatory functions quantitatively

the setup:




assumptions:

- fluorescence ~ concentration
- well isolated system


need to:

- calibrate expression
- quantify parameters
- make sure degradation is negligible
- measure cl and gene expression simultaneously

checks:





results:






Sequestering of TFs in complexes



- yeast
- synthetic circuit - but common motif proteins
- 1 binding site (no cooperativity)
- measure active TF and gene expression (YFP)
- also achieved by small RNAs, enzymes, ...



→ ultrasensitivity



no sequestration




with sequestration

The lac operon

Function: break up **lactose**

Jacob, Monod, Lwoff and collaborators (1950s)



....
 presentation taken from Terry Hwa

lacZ - encodes intracellular enzyme that cleaves lactose into glucose and galactose

lacY - encodes a membrane-bound transport protein that pumps lactose into the cell

lacA- encodes another enzyme

Expressed only in the **presence** of **lactose** and **absence** of **glucose**

- repressed by the **Lac Repressor** (encoded by *lacI*)
- repression alleviated by allo-lactose (by-product of lactose metabolism)
- synthetically allo-lactose replaced by **IPTG**
- activated by the global regulator **Crp**; requires the inducer **cAMP**
- cAMP synthesized endogenously by **Adenylate Cyclase** (encoded by *cyaA*)
- activity of **AC** repressed by **glucose** uptake

IPTG	glucose	expression
low	high	OFF
low	low	OFF
high	high	OFF
high	low	ON

1960's






Fig.1. Growth of *Escherichia coli* in the presence of different carbohydrate pairs serving as the only source of carbon in a synthetic medium¹⁰.




[Jacob, Monod]


2000's



microarrays




fluorescent proteins




[Ozbudak et al]

The lac operon



molecular complexity:

- protein-protein interactions
- specific protein-DNA binding
- looping



Y. Setty, A.E. Mayo, M.G. Surette, U. Alon
Detailed map of a cis-regulatory input function
PNAS June 24, 2003 vol. 100 no. 13 7702

Initial experiments:


- grow cells in medium with **cAMP** and **IPTG**
- use **glucose** to suppress intracellular **cAMP** synthesis
- control **cAMP**-level extracellularly

Problem:

- inconsistency with theory of mutants


Possible reason:

- extracellular **cAMP** and **IPTG** concentrations (the ones that are controlled) not equal to intracellular ones (the ones that matter)




The lac operon

theory



E. coli MG1655
(*cyaA*⁻, *cpdA*⁻, *lacY*⁻)




Solutions of problem:


- the sensitivity of the inducer response results from LacI-mediated DNA looping
- DNA looping is significantly enhanced by **Crp**.
- **IPTG** intake increased by LacY
- difference between **IPTG** and **lactose** uptake mechanisms

Lesson:

- quantitative study leads to correct mechanistic understanding







Cooperativity




Monod-Wyman-Changeux (MWC) model

Post-translational regulation






Yeast protein-protein interaction network





Yeast protein interaction network: 1870 proteins, 2240 interactions

H. Jeong et al, Nature 411,41 (2001)

Relative proportion of different FFL, Mangan et al J. Mol. Biol (2006)




Coherent feed-forward loop




Alon, Nat Rev Gen (2007)


b Arabinose system




Lac system




ON step of S_x



OFF step of S_x





Alon, Nat Rev Gen (2007)



Alon, Nat Rev Gen (2007)

Incoherent feed-forward loop or feed-forward inhibition




Alon, Nat Rev Gen (2007)

The λ phage

XOR promoter!

looping
dimer binding





Lysogeny









cro - low
cl - high

Lysis

cro - high
cl - low





The synthetic toggle switch




Timothy S. Gardner, Charles R. Cantor & James J. Collins
Construction of a genetic toggle switch in *Escherichia coli*


$$\frac{du}{dt} = \frac{\alpha_1}{1+v^\beta} - u$$

$$\frac{dv}{dt} = \frac{\alpha_2}{1+u^\gamma} - v$$






Hirata et al, Nat. Gen. 36, 750 (2004)






The repressilator



$$\frac{dm_i}{dt} = -m_i + \frac{\alpha}{(1 + p_i^n)} + \alpha_0 \quad (i = lacI, tetR, cl)$$

$$\frac{dp_j}{dt} = -\beta(p_j - m_j) \quad (j = cl, lacI, tetR)$$




Michael B. Elowitz & Stanislas Leibler



A synthetic oscillatory network of transcriptional regulators

NATURE | VOL 403 | 20 JANUARY 2000





Predator-prey oscillations



$$\begin{cases} \frac{d[A]}{dt} = \alpha \cdot \mathcal{G}_A([A]) \cdot \mathcal{G}_R([R]) - \beta \cdot [A] \\ \frac{d[R]}{dt} = \alpha \cdot \mathcal{G}_A([A]) - \beta \cdot [R] \end{cases}$$



Predator-prey oscillations















Mariette R. Atkinson, Michael A. Savageau, Jesse T. Myers,
and Alexander J. Ninfa

Development of Genetic Circuitry Exhibiting Toggle Switch or
Oscillatory Behavior in *Escherichia coli*

Cell, Vol. 113, 597-607, May 30, 2003

Predator-prey oscillations



Jesse Stricker, Scott Cookson, Matthew R. Bennett, William H. Mather, Lev S. Tsimring & Jeff Hasty

A fast, robust and tunable synthetic gene oscillator

NATURE Vol 456|27 November 2008

$$\frac{d[\text{Cdc13}_T]}{dt} = k_1 M - \left(k'_2 + k''_2 [\text{Ste9}] + k'''_2 [\text{Slp1}] \right) [\text{Cdc13}_T],$$

$$\frac{d[\text{preMPF}]}{dt} = k_{\text{wee}} ([\text{Cdc13}_T] - [\text{preMPF}]) - k_{25} [\text{preMPF}] - \left(k'_2 + k''_2 [\text{Ste9}] + k'''_2 [\text{Slp1}] \right) [\text{preMPF}],$$

$$\frac{d[\text{Ste9}]}{dt} = \left(k'_3 + k''_3 [\text{Slp1}] \right) \frac{1 - [\text{Ste9}]}{J_3 + 1 - [\text{Ste9}]} - \left(k'_4 [\text{SK}] + k_4 [\text{MPF}] \right) \frac{[\text{Ste9}]}{J_4 + [\text{Ste9}]},$$

$$\frac{d[\text{Slp1}]}{dt} = k'_5 + k''_5 \frac{[\text{MPF}]^4}{J_5^4 + [\text{MPF}]^4} - k_6 [\text{Slp1}_T],$$

$$\frac{d[\text{Slp1}^*]}{dt} = k_7 [\text{IEP}] \frac{[\text{Slp1}_T] - [\text{Slp1}]}{J_7 + [\text{Slp1}_T] - [\text{Slp1}]} - k_8 \frac{[\text{Slp1}]}{J_8 + [\text{Slp1}]} - k_6 [\text{Slp1}],$$

$$\frac{d[\text{IEP}]}{dt} = k_9 [\text{MPF}] \frac{1 - [\text{IEP}]}{J_9 + 1 - [\text{IEP}]} - k_{10} \frac{[\text{IEP}]}{J_{10} + [\text{IEP}]},$$

$$k_{\text{wee}} = k'_{\text{wee}} + \left(k''_{\text{wee}} - k'_{\text{wee}} \right) G(V_{\text{awee}}, V_{\text{iwee}} [\text{MPF}], J_{\text{awee}}, J_{\text{iwee}}),$$

$$k_{25} = k'_{25} + \left(k''_{25} - k'_{25} \right) G(V_{\text{a25}} [\text{MPF}], V_{\text{i25}}, J_{\text{a25}}, J_{\text{i25}}),$$

$$\Sigma = [\text{Cdc13}_T] + [\text{Rum1}_T] + K_{\text{diss}}.$$

$$G(a, b, c, d) = \frac{2ad}{b - a + bc + ad + \sqrt{(b - a + bc + ad)^2 - 4ad(b - a)}}$$

$$\frac{d[\text{Rum1}_T]}{dt} = k_{11} - \left(k_{12} + k'_{12} [\text{SK}] + k''_{12} [\text{MPF}] \right) [\text{Rum1}_T],$$

$$\frac{d[\text{SK}]}{dt} = k_{13} [\text{TF}] - k_{14} [\text{SK}],$$

$$\frac{d[M]}{dt} = \mu M,$$

$$[\text{Trimer}] = \frac{2[\text{Cdc13}_T][\text{Rum1}_T]}{\Sigma + \sqrt{\Sigma^2 - 4[\text{Cdc13}_T][\text{Rum1}_T]}},$$

$$[\text{MPF}] = \frac{([\text{Cdc13}_T] - [\text{preMPF}])([\text{Cdc13}_T] - [\text{Trimer}])}{[\text{Cdc13}_T]},$$

$$[\text{TF}] = G(k_{15}M, k'_{16} + k''_{16}[\text{MPF}], J_{15}, J_{16})$$

- 9 ODEs
- 3 algebraic equations
- ~30 parameters

Novak et al, Chaos 11(1):277–286, 2001