

Symmetry resolved entanglement



Pasquale Calabrese
SISSA-Trieste




Hubert's fest, Sept 2021



Reduced density matrix and entanglement

Consider a system in a quantum state $|\psi\rangle$, $\rho = |\psi\rangle\langle\psi|$

$$\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B \quad |\psi\rangle \neq |\psi_A\rangle \otimes |\psi_B\rangle$$



The reduced density matrix of A is $\rho_A = \text{Tr}_B |\psi\rangle\langle\psi|$

The entanglement entropy

$$S_A = -\text{Tr}(\rho_A \log \rho_A)$$

measures the **bipartite entanglement** between A & B


We will also use the Rènyi entropies $S_n = \frac{1}{1-n} \log \text{Tr} \rho_A^n$

Entanglement and symmetries

Let's assume that $|\psi\rangle$ is symmetric under the action of a charge Q , i.e $[\rho, Q] = 0$

The charge is local: $Q = Q_A + Q_B$

$$[\rho, Q] = 0 \xrightarrow{\text{Tr}_B} [\rho_A, Q_A] = 0$$


 ρ_A has a block diagonal form:

$$\rho_A = \bigoplus_q \Pi_q \rho_A = \bigoplus_q [p(q) \rho_A(q)] \quad \text{with} \quad p(q) = \text{Tr}(\Pi_q \rho_A)$$

Symmetry resolved entanglement entropy:

$$S(q) = -\text{Tr}[\rho_A(q) \log \rho_A(q)]$$

$$\rho_A = \begin{pmatrix} \boxed{q_1} & & & \\ & \boxed{q_2} & & \\ & & \boxed{q_3} & \\ & & & \ddots \end{pmatrix}$$

 probability of being in the sector q

Entanglement and symmetries II

The symmetry resolved entanglement satisfies the sum rule

$$S = \sum_q p(q) S(q) - \sum_q p(q) \log(p(q)) \equiv S^c + S^n$$

S^c : Configurational entropy

S^n : Number entropy





FIG. S8. **Total entropy partitioned** The total von Neumann entanglement entropy S_{vN} for the half-system is shown as a function of time in an interacting system at strong disorder. The entropy is split up into S_n and S_c . For visual



A. Lukin, M. Rispoli, R. Schittko, M. E. Tai, A. M. Kaufman, S. Choi, V. Khemani, J. Leonard, and M. Greiner, Probing entanglement in a many-body localized system, Science 364, 6437 (2019).

Entanglement and symmetries: results

Early work

N. Laflorencie and S. Rachel, J. Stat. Mech. (2014) P11013

SRE in CFT

M. Goldstein and E. Sela, PRL 120, 200602 (2018)
J.C. Xavier, F.C. Alcaraz, and G. Sierra, PRB 98, 041106 (2018)
L. Capizzi, P. Ruggiero, and P. Calabrese, JSTAT (2020) 073101
R. Bonsignori and P. Calabrese, JPA 54, 015005 (2020)
B. Estienne et al, SciPost Phys. 10, 54 (2021)
S. Murciano, J. Dubail, P. Calabrese, ArXiv 2106.15946

Relative entropy and distances:

H.-H. Chen, arXiv:2104.03102
L. Capizzi and P. Calabrese, ArXiv:2105.08596

Free QFT

S. Murciano, G. Di Giulio, and P. Calabrese, JHEP 08 (2020) 073

Holography

S. Zhao, C. Northe, and R. Meyer, arXiv:2012.11274

Disorder Systems

X. Turkeshi, P. Ruggiero, V. Alba,
and P. Calabrese, PRB 102, 014455 (2020)

Negativity

E. Cornfeld, M. Goldstein, and E. Sela, PRA 98, 032302 (2018)
S. Murciano, R. Bonsignori, P. Calabrese, SciPost Phys. 10, 111 (2021)
P. Calabrese, P. Zoller, B. Vermersch, R. Kueng,
and B. Kraus, ArXiv:2103.07443

Lattice free fermions

R. Bonsignori, P. Ruggiero, and P. Calabrese, JPA 52, 475302 (2019)
M. T. Tan and S. Ryu, PRB 101, 235169 (2020)
S. Fraenkel and M. Goldstein, JSTAT 033106 (2020)
S. Murciano, P. Ruggiero, and P. Calabrese, JSTAT (2020) 083102

Non-equilibrium and quantum quenches

N. Feldman and M. Goldstein, PRB 100, 235146 (2019)
G. Perez, R. Bonsignori and P. Calabrese, PRB 103, L041104 (2020)
V. Vitale, A. Elben, R. Kueng, A. Neven, J. Carrasco, B. Kraus, P. Zoller,
P. Calabrese, B. Vermersch, and M. Dalmonte, ArXiv:2101.07814
S. Fraenkel and M. Goldstein, ArXiv:2105.00740
G. Perez, R. Bonsignori and P. Calabrese, ArXiv:2106.13115

Integrability

Corner Transfer Matrix

S. Murciano, G. Di Giulio, and P. Calabrese, SciPost Phys. 8, 046 (2020)
P. Calabrese, M. Collura, G. Di Giulio, and S. Murciano, EPL 129, 60007 (2020)

Form Factor Bootstrap

D. X. Horvath and P. Calabrese, JHEP 11 131 (2020)
D. X. Horvath, L. Capizzi, and P. Calabrese, JHEP 05 197 (2021)
D. X. Horvath, P. Calabrese, and O. A. Castro-Alvaredo, arXiv:2105.13982
L. Capizzi, D. Horvath, P. Calabrese, and O. Castro-Alvaredo arXiv:2108.10935

Topology

E. Cornfeld, L. A. Landau, K. Shtengel, and E. Sela, PRB 99, 115429 (2019)
K. Monkman and J. Sirker, arXiv:2005.13026
D. Azses and E. Sela, arXiv:2008.09332

Entanglement and symmetries: results

Early work

N. Laflorencie and S. Rachel, J. Stat. Mech. (2014) P11013

SRE in CFT

- M. Goldstein and E. Sela, PRL 120, 200602 (2018)
- J. S. Xavier, F.C. Alcaraz, and G. Sierra, PRB 98, 041106 (2018)
- L. Capizzi, P. Ruggiero, and P. Calabrese, JSTAT (2020) 073101
- R. Bonsignori and P. Calabrese, JPA 54, 015005 (2020)
- B. Estienne et al, SciPost Phys. 10, 54 (2021)
- S. Murciano, J. Dubail, P. Calabrese, ArXiv 2106.15946

Relative entropy and distances:

H.-H. Chen, arXiv:2104.03102
L. Capizzi and P. Calabrese, ArXiv:2105.08596

Free QFT

S. Murciano, G. Di Giulio, and P. Calabrese, JHEP 08 (2020) 073

Disorder Systems

X. Turkeshi, P. Ruggiero, V. Alba,
and P. Calabrese, PRB 102, 014455 (2020)

Holography

S. Zhao, C. Northe, and R. Meyer, arXiv:2012.11274

Negativity

- E. Cornfeld, M. Goldstein, and E. Sela, PRA 98, 032302 (2018)
- S. Murciano, R. Bonsignori, P. Calabrese, SciPost Phys. 10, 111 (2021)
- P. Calabrese, P. Zoller, B. Vermersch, R. Kueng,
and B. Kraus, ArXiv:2103.07443

Lattice free fermions

- R. Bonsignori, P. Ruggiero, and P. Calabrese, JPA 52, 475302 (2019)
- M. T. Tan and S. Ryu, PRB 101, 235169 (2020)
- S. Fraenkel and M. Goldstein, JSTAT 033106 (2020)
- S. Murciano, P. Ruggiero, and P. Calabrese, JSTAT (2020) 083102

Integrability

Corner Transfer Matrix

S. Murciano, G. Di Giulio, and P. Calabrese, SciPost Phys. 8, 046 (2020)
P. Calabrese, M. Collura, G. Di Giulio, and S. Murciano, EPL 129, 60007 (2020)

Form Factor Bootstrap

D. X. Horvath and P. Calabrese, JHEP 11 131 (2020)
D. X. Horvath, L. Capizzi, and P. Calabrese, JHEP 05 197 (2021)
D. X. Horvath, P. Calabrese, and O. A. Castro-Alvaredo, arXiv:2105.13982
L. Capizzi, D. Horvath, P. Calabrese, and O. Castro-Alvaredo arXiv:2108.10935

Non-equilibrium and quantum quenches

N. Feldman and M. Goldstein, PRB 100, 235146 (2019)
G. Perez, R. Bonsignori and P. Calabrese, PRB 103, L041104 (2020)
V. Vitale, A. Elben, R. Kueng, A. Neven, J. Carrasco, B. Kraus, P. Zoller,
P. Calabrese, B. Vermersch, and M. Dalmonte, ArXiv:2101.07814
S. Fraenkel and M. Goldstein, ArXiv:2105.00740
G. Perez, R. Bonsignori and P. Calabrese, ArXiv:2106.13115


Topology

E. Cornfeld, L. A. Landau, K. Shtengel, and E. Sela, PRB 99, 115429 (2019)
K. Monkman and J. Sirker, arXiv:2005.13026
D. Azses and E. Sela, arXiv:2008.09332

Entanglement entropy and path integral

PC, J Cardy 2004

The density matrix at temperature $1/\beta$ is


$$\langle \Phi_1 | \rho | \Phi_2 \rangle =$$


$$= \int \frac{[d\phi(x, \tau)]}{Z} \prod_x \delta(\phi(x, 0) - \phi_2(x)) \prod_x \delta(\phi(x, \beta) - \phi_1(x)) e^{-S_E}$$

The trace sews together the edges along $\tau = 0$ and $\tau = \beta$ to form a cylinder of circumference β .

$A = (u, v)$: ρ_A sews together only those points x which are not in A , leaving an open cut at $\tau = 0$

$$\langle \Phi_1(x) | \rho_A | \Phi_2(x) \rangle =$$



Replicas and Riemann surfaces


PC, J Cardy 2004

$$S_A = -\text{Tr}(\rho_A \log \rho_A) = -\lim_{n \rightarrow 1} \frac{\partial}{\partial n} \text{Tr}(\rho_A^n)$$

For n **integer**, $\text{Tr} \rho_A^n$ is obtained by sewing cyclically n cylinders above.

This is the partition function on a **n -sheeted Riemann surface**

$$\text{Tr} \rho_A^n =$$




$$\text{Renyi entanglement entropies } S_n = \frac{1}{1-n} \text{Tr} \rho_A^n$$

Riemann surfaces and CFT


This Riemann surface is mapped to the plane by

PC, J Cardy 2004

$$w \rightarrow \zeta = \frac{w-u}{w-v}; \zeta \rightarrow z = \zeta^{1/n} \Rightarrow w \rightarrow z = \left(\frac{w-u}{w-v} \right)^{1/n}$$



$$\text{Tr} \rho_A^n =$$



$$= c_n |u - v|^{-\frac{c}{6}(n-1/n)} \xrightarrow{|u-v|=\ell} S_A = -\lim_{n \rightarrow 1} \frac{\partial}{\partial n} \text{Tr}(\rho_A^n) = \frac{c}{3} \log \ell$$

$\text{Tr} \rho_A^n$ is equivalent to the 2-point function of **twist fields**

$$\text{Tr} \rho_A^n = \langle \mathcal{T}_n(u) \bar{\mathcal{T}}_n(v) \rangle \quad \text{with scaling dimension}$$

$$\Delta_{\mathcal{T}_n} = \frac{c}{12} \left(n - \frac{1}{n} \right)$$

U(1) Symmetry resolution in CFT

M. Goldstein and E. Sela, PRL 120, 200602 (2018)




Symmetry resolved Renyi: $S_n(q) \equiv \frac{1}{1-n} \log \text{Tr} \rho_A^n(q)$


It requires the resolution of the spectrum in Q

Introduce the charged moments:

$$Z_n(\alpha) \equiv \text{Tr} \rho_A^n e^{iQ_A \alpha}$$



$$\mathcal{Z}_n(q) \equiv \text{Tr}(\Pi_q \rho_A^n) = \int_{-\pi}^{\pi} \frac{d\alpha}{2\pi} e^{-iq\alpha} Z_n(\alpha)$$




$$S_n(q) = \frac{1}{1-n} \log \left[\frac{\mathcal{Z}_n(q)}{\mathcal{Z}_1(q)^n} \right], \quad S_1(q) = -\partial_n \left[\frac{\mathcal{Z}_n(q)}{\mathcal{Z}_1(q)^n} \right]_{n=1}$$

$Z_n(\alpha)$ is the partition function in the presence of a charge flux.


The field takes a total phase α going through $\mathcal{R}_n$.

In CFT it is placed all in one sheet introducing the composite field:


$$\mathcal{T}_{n,\alpha}(x, \tau) \phi_i(x', \tau) = \begin{cases} \phi_{i+1}(x', \tau) e^{i\alpha \delta_{i,n}} \mathcal{T}_{n,\alpha}(x, \tau) & (x < x'), \\ \phi_i(x', \tau) \mathcal{T}_{n,\alpha}(x, \tau) & \text{otherwise.} \end{cases}$$




$$p(q) = \mathcal{Z}_1(q)$$



$$Z_n(\alpha) = \langle \mathcal{T}_{n,\alpha}(\ell, 0) \tilde{\mathcal{T}}_{n,\alpha}(0, 0) \rangle$$



$$h_{n,\alpha} = h_n + \frac{h_\alpha}{n}$$



$$Z_n(\alpha) = c_{n,\alpha} \ell^{-\frac{c}{6} \left(n - \frac{1}{n} \right) - 2 \frac{h_\alpha + \bar{h}_\alpha}{n}}$$

Symmetry resolution in CFT: compact boson

Action: $\mathcal{S}_E[\varphi] = \frac{1}{8\pi K} \int d\tau dx \partial_\mu \varphi \partial^\mu \varphi$ Conserved charge $Q_A = \frac{1}{2\pi} \int_A \partial \varphi(x,0) dx \Rightarrow e^{i\alpha Q_A} = e^{i\frac{\alpha}{2\pi} \varphi(u,0)} e^{-i\frac{\alpha}{2\pi} \varphi(v,0)}$

$\Rightarrow h_\alpha = \bar{h}_\alpha = \frac{1}{2} \left(\frac{\alpha}{2\pi} \right)^2 K \Rightarrow Z_n(\alpha) = c_{n,\alpha} \ell^{-\frac{c}{6}(n-\frac{1}{n}) - \frac{2K}{n} \left(\frac{\alpha}{2\pi} \right)^2}$

Fourier transform using saddle point

$\Rightarrow \mathcal{Z}_n(q) = c_n \ell^{-\frac{c}{6}(n-\frac{1}{n})} \sqrt{\frac{n\pi}{2K \ln \ell + \gamma_n}} e^{\frac{n\pi^2(q - \langle Q_A \rangle)^2}{2K \ln \ell + \gamma_n}}$

$\Rightarrow S_n(q) = S_n - \frac{1}{2} \log \left(\frac{2K}{\pi} \log \ell \right) + \frac{\log n}{2(1-n)} + o(\ell^0)$

Entanglement equipartition: up to order $o(1)$, the SR entanglement does not depend on the symmetry sector

Symmetry resolution in CFT: compact boson II

R. Bonsignori, P. Ruggiero, and P. Calabrese, JPA 52, 475302 (2019)

$$S_n(q) = S_n - \frac{1}{2} \log \left(\frac{2K}{\pi} \log \ell \right) + \frac{\log n}{2(1-n)} + o(\ell^0)$$

Q: Where the log log term ends up in the total entropy?

A: It is exactly canceled by the number entropy:

$$S = \sum_q p(q) S(q) - \sum_q p(q) \log(p(q)) \equiv S^c + S^n$$

$$S^n = \frac{1}{2} + \frac{1}{2} \ln \left(\frac{2K}{\pi} \ln \ell \right) + o(1)$$

Note: The number entropy satisfies $S^n \ll S \sim S(q)$, a fact valid much more generally

Lattice free fermions



R. Bonsignori, P. Ruggiero, and P. Calabrese, JPA 52, 475302 (2019)

$$H = - \sum_{i=-\infty}^{\infty} \left[c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i - 2h \left(c_i^\dagger c_i - \frac{1}{2} \right) \right]$$

Using Fisher Hartwig techniques:

$$\ln Z_n^{(0)}(\alpha) = i\alpha \frac{k_F \ell}{\pi} - \left[\frac{1}{6} \left(n - \frac{1}{n} \right) + \frac{2}{n} \left(\frac{\alpha}{2\pi} \right)^2 \right] \ln L_k + \Upsilon(n, \alpha)$$




$$\Upsilon(n, \alpha) = ni \int_{-\infty}^{\infty} dw [\tanh(\pi w) - \tanh(\pi n w + i\alpha/2)] \ln \frac{\Gamma(\frac{1}{2} + iw)}{\Gamma(\frac{1}{2} - iw)}$$



Fourier tranform + ratios for entropies



$$S_n(q) = S_n - \frac{1}{2} \ln \left(\frac{2}{\pi} \ln \delta_n L_k \right) + \frac{\ln n}{2(1-n)} + (q - \bar{q})^2 \pi^4 \frac{n(\gamma_2(1) - n\gamma_2(n))}{1-n} \frac{1}{\ln^2 \kappa_n L_k} + \dots$$

Equipartition is broken at order $(\log \ell)^{-2}$




2D free lattice QFT: dimensional reduction

S. Murciano, P. Ruggiero, and P. Calabrese, JSTAT (2020) 083102




Idea: reduce the initial 2D system into decoupled 1D ones in a mixed space-momentum representation

Fermions



Bosons



Resolution of non-abelian symmetries: WZW models

S. Murciano, J. Dubail, P. Calabrese, ArXiv:2106.15946

Consider a general non-abelian group G (of dimension d and volume $\text{Vol}(G)$) and the corresponding WZW model

$$\rho_A = \bigoplus_r [p(r)\rho_A(r)] \quad r \text{ labels the irreducible representations of } G, \dim(r) \text{ its dimension}$$

$SU(2)$ done by Goldstein and Sela in 2018 paper using $SU(2)$ algebra

Our Strategy (without mentioning many highly non trivial points and assumptions)

- Write the charged moments as a linear combination of the unspecialised characters
- Use their modular properties to compute the resolved partition functions, by identifying all states in a given representation of the group.
- The SR entropies are obtained integrating the group characters around all saddles (that are the elements of the center $Z(G)$ (of order $|Z(G)|$))

Final Result

$$S_n^r(L) = S_n(L) - \frac{d}{2} \log(\log L) + 2 \log \dim(r) - \log \frac{\text{Vol}(G)}{|Z(G)|} + \frac{d}{2} \left(-\log k + \frac{\log n}{1-n} + \log(2\pi^3) \right) + o(L^0)$$


Equipartition broken at order $O(1)!!$

SRE after a quantum quench

Prepare a system in a low-entangled initial state $|\psi_0\rangle$ and let it evolve unitarily $|\psi(t)\rangle = e^{iHt}|\psi_0\rangle$

Long story short: In integrable models the entanglement dynamics is captured by the **quasiparticle picture**

PC & Cardy, 2005 + Alba & PC 2017



$$S = \int \frac{dk}{2\pi} h(k) \min[2v_k t, \ell]$$

Adapting the QP picture to the charged moments, we conjecture for a general integrable model

G. Perez, R. Bonsignori and P. Calabrese, PRB 103, L041104 (2020)

$$\log Z_n(\alpha) = i\langle Q_A \rangle \alpha + \int \frac{dk}{2\pi} f_{n,\alpha}(k) \min[2v_k t, \ell],$$

but the kernel $f_{n,\alpha}(k)$ is difficult to compute for generic model, while free is possible




FIG. 1. The time evolution of the charged moments $Z_n(\alpha)$ after a quench from the Néel state in the free fermion model

SRE after a quantum quench II

G. Perez, R. Bonsignori and P. Calabrese, PRB 103, L041104 (2020)

Some general features in charge space:

Delay time $t_D \propto |\Delta q|$

$$t_D = \pi \frac{|\Delta q|}{4} \quad \text{for free fermions}$$



The time needed to change the charge of an amount $|\Delta q|$ within A

Equipartition for small $|\Delta q|$

$$S_n(q) = S_n - \frac{\Delta q^2}{4(1-n)} \left\{ \frac{1}{\mathcal{J}_n} - \frac{n}{\mathcal{J}_1} \right\}$$

Number entropy

$$S^n \simeq \frac{1}{2} \log t$$





Application to ion-trap experiment: SR dynamical purification

V. Vitale, A. Elben, R. Kueng, A. Neven, J. Carrasco, B. Kraus, P. Zoller, P. Calabrese, B. Vermersch, and M. Dalmonte, ArXiv:2101.07814

Hamiltonian + dissipative dynamics
$$\partial_t \rho = -\frac{i}{\hbar} [H, \rho] + \sum_j \gamma \left[b_j \rho b_j^\dagger + b_j^\dagger \rho b_j - \frac{1}{2} \{b_j b_j^\dagger + n_j, \rho\} \right]$$



- Recap:**
- Both dynamics leads to entropy growth (entanglement and total)
 - The total entropy grows, purity reduces

Analysis of experimental results:




Some sectors purifies at intermediate times

A general phenomenon that can be easily shown in perturbation theory in γ



Mixed state entanglement: Partial transpose and negativity

Q: what is the entanglement in a mixed state?



$|e_k^1\rangle$ and $|e_l^2\rangle$ bases of A_1 and A_2

$$\rho_A = \sum_{ijkl} \langle e_i^1, e_j^2 | \rho_A | e_k^1, e_l^2 \rangle |e_i^1, e_j^2\rangle \langle e_k^1, e_l^2|$$



$$\rho_A^{T_1} = \sum_{ijkl} \langle e_k^1, e_j^2 | \rho_A | e_i^1, e_l^2 \rangle |e_k^1, e_j^2\rangle \langle e_i^1, e_l^2|$$


$$(|e_i^1, e_j^2\rangle \langle e_k^1, e_l^2|)^{T_1} \equiv |e_k^1, e_j^2\rangle \langle e_i^1, e_l^2|$$



PPT criterion:

If $\rho_A^{T_1}$ has negative eigenvalues
 ρ_A is entangled

Peres, 1996



The Negativity $= \mathcal{N} = \frac{\text{Tr} |\rho_A^{T_1}| - 1}{2}$ measure how much the eigenvalues of $\rho_A^{T_1}$ are negative and it is an entanglement monotone

Vidal Werner 2002

Replica trick: $\text{Tr} |\rho_A^{T_1}| = \lim_{n \rightarrow 1/2} \text{Tr}(\rho_A^{T_1})^{2n}$

P. Calabrese, J. Cardy, E. Tonni 2012

Intermezzo: “Negativity” in experiments

The negativity is difficult to measure experimentally, but the moments of the partial transpose p_n can

E. Cornfeld, M. Goldstein, and E. Sela, PRA 98, 032302 (2018)

J. Gray, L. Banchi, A. Bayat, and S. Bose, Phys. Rev. Lett. 121, 150503 (2018)

A. Elben, R. Kueng, H.-Y. Huang, R. van Bijnen, C. Kokail, M. Dalmonte, P. Calabrese, B. Kraus, J. Preskill, P. Zoller, and B. Vermersch, PRL 125, 200501 (2020)



In Elben et al, PRL 2020 p_n are obtained by performing local random measurements and post-processing using the classical shadows framework

p_3 -PPT condition: if $p_3 < p_2^2$, then PPT is violated and there is entanglement

Generalizations

A. Neven, J. Carrasco, V. Vitale, C. Kokail, A. Elben, M. Dalmonte, P. Calabrese, P. Zoller, B. Vermersch, R. Kueng, and B. Kraus, ArXiv:2103.07443


- D_n conditions: generalized conditions, involving higher moments
- Symmetry resolution of **p_3 -PPT**:
 - Allow to understand in which sector negative eigenvalues are
 - More sensitive to small negative eigenvalues



Fermionic partial transpose

H. Shapourian, K. Shiozaki, S. Ryu, PRB 95, 165101 (2017)


Occupation number basis: $|\{n_j\}_{j \in A_1}, \{n_j\}_{j \in A_2}\rangle = (f_{m_1}^\dagger)^{n_{m_1}} \dots (f_{m_{\ell_1}}^\dagger)^{n_{m_{\ell_1}}} (f_{m'_1}^\dagger)^{n_{m'_1}} \dots (f_{m'_{\ell_2}}^\dagger)^{n_{m'_{\ell_2}}} |0\rangle$



$\rho_A^{R_1}$ non hermitian

Fermionic partial transpose:

$$(|\{n_j\}_{A_1}, \{n_j\}_{A_2}\rangle \langle \{\bar{n}_j\}_{A_1}, \{\bar{n}_j\}_{A_2}|)^{R_1} = (-1)^{\phi(\{n_j\}, \{\bar{n}_j\})} (|\{\bar{n}_j\}_{A_1}, \{n_j\}_{A_2}\rangle \langle \{n_j\}_{A_1}, \{\bar{n}_j\}_{A_2}|)$$



$$\mathcal{N} = \frac{\text{Tr} |\rho_A^{R_1}| - 1}{2} = \frac{\text{Tr} \sqrt{\rho_A^{R_1} (\rho_A^{R_1})^\dagger} - 1}{2} = \lim_{n \rightarrow 1/2} \frac{\text{Tr} (\rho_A^{R_1} (\rho_A^{R_1})^\dagger)^n - 1}{2}$$

Fermionic negativity (no negative eigenvalues, but entanglement monotone)

$$\rho_A^{T_1} = \frac{e^{i\pi/4} \rho_A^{R_1} + e^{-i\pi/4} (\rho_A^{R_1})^\dagger}{\sqrt{2}}$$

$\text{Tr}(\rho_A^{T_1})^{2n}$
 $\sum$ all spin structures
 $\rho_A^{T_1}$ sum of 2 Gaussian

$\text{Tr}(\rho_A^{R_1} (\rho_A^{R_1})^\dagger)^n$
 only 1 cycle
 $\rho_A^{R_1}$ Gaussian

Symmetry Resolution: example

a particle in one out of three boxes, $A = A_1 \cup A_2, B,$

$$|\Psi\rangle = \alpha |100\rangle + \beta |010\rangle + \gamma |001\rangle$$

$$\rho_A = \begin{pmatrix} |00\rangle & |01\rangle & |10\rangle & |11\rangle \\ \begin{pmatrix} |\gamma|^2 & 0 & 0 & 0 \end{pmatrix} \\ \begin{pmatrix} 0 & |\beta|^2 & \alpha^* \beta \\ 0 & \beta^* \alpha & |\alpha|^2 \end{pmatrix} \\ \begin{pmatrix} 0 & 0 & 0 & 0 \end{pmatrix} \end{pmatrix}$$

block-diagonal structure

(fermionic)
partial transpose

$$\rho_A^{R_1} = \begin{pmatrix} |\gamma|^2 & 0 & \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \begin{pmatrix} i\alpha\beta^* \\ 0 \end{pmatrix} \\ 0 & |\beta|^2 & \begin{pmatrix} 0 \\ i\beta\alpha^* \end{pmatrix} & \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ \begin{pmatrix} 0 \\ i\beta\alpha^* \end{pmatrix} & \begin{pmatrix} 0 \\ 0 \end{pmatrix} & |\alpha|^2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

block-diagonal structure

$$\rho_A^{R_1} \cong (|\alpha|^2)_{q=-1} \oplus \begin{pmatrix} |\gamma|^2 & i\alpha\beta^* \\ i\beta\alpha^* & 0 \end{pmatrix}_{q=0} \oplus (|\beta|^2)_{q=1}$$

$$Q = Q_{A_2} - Q_{A_1}^{R_1}$$

charge imbalance resolved negativity

$$\rho_A \cong (|\gamma|^2)_{\tilde{q}=0} \oplus \begin{pmatrix} |\beta|^2 & \alpha\beta^* \\ \beta\alpha^* & |\alpha|^2 \end{pmatrix}_{\tilde{q}=1} \oplus (0)_{\tilde{q}=2}$$

$$\mathcal{N}(q) = \frac{\text{Tr} |(\rho_A^{R_1}(q))| - 1}{2}, \quad \rho_A^{R_1}(q) = \frac{\mathcal{P}_q \rho_A^{R_1} \mathcal{P}_q}{\text{Tr}(\mathcal{P}_q \rho_A^{R_1})}$$







E. Cornfeld, M. Goldstein, and E. Sela, PRA 98, 032302 (2018)

$$\mathcal{N} = \sum_q p(q) \mathcal{N}(q), \quad p(q) = \text{Tr}(\mathcal{P}_q \rho_A^{R_1})$$

S. Murciano, R. Bonsignori, P. Calabrese, SciPost Phys 10, 111(2021)


Some results: Negativity equipartition

S. Murciano, R. Bonsignori, P. Calabrese, SciPost Phys 10, 111(2021)



Main message:

The symmetry resolution of entanglement measures provides a fine structure of the entanglement content of physical states of extended quantum systems that is not accessible from the measure of the total entanglement



Some features:

- Measurable experimentally (actually already measured!)
- Easy to compute via charged moments
- Relation to charge statistics, entanglement Hamiltonian,

HAPPY B'DAY HUBERT!