Elasticity of a contact-line and avalanche-size distribution at depinning

Pierre Le Doussal and Kay Jörg Wiese

CNRS-Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, 73231 Paris Cedex 05, France

August 27, 2009

Abstract. Motivated by recent experiments, we extend the Joanny-deGennes calculation of the elasticity of a contact line to an arbitrary contact angle and an arbitrary plate inclination in presence of gravity. This requires a diagonalization of the elastic modes around the non-linear equilibrium profile, which is carried out exactly. We then make detailed predictions for the avalanche-size distribution at quasi-static depinning: we study how the universal (i.e. short-scale independent) rescaled size distribution and the ratio of moments of local to global avalanches depend on the precise form of the elastic kernel.

PACS. 68.35.Rh Phase transitions and critical phenomena

1 Introduction

Recent experiments on contact-line dynamics near depinning [1] have allowed for a direct measurement of the renormalized disorder correlator, and of the avalanche-size statistics. It was concluded in [1] that the precise form of the contact-line elasticity may have an important impact on the distribution of avalanche sizes. Thus, a detailed prediction for the former is needed. Surprisingly, a review of the literature reveals that this problem appears to have been treated only neglecting gravity [2]¹, or in presence of gravity only in the case where the equilibrium configuration of the interface is flat and horizontal, as for a vertical wall and a contact angle θ of 90° [3,4], see Figs. 1 and 2.

¹ [2] introduce the capillary length L_c ad hoc as a cutoff for the contact-line profile in response to a δ -like force.



Fig. 1. The coordinate system for a vertical wall. The air/liquid interface becomes flat for $x \gg L_c$.



Fig. 2. The coordinate system for a wall inclined by φ with respect to the vertical: the coordinate y, not shown, is perpendicular to the plane of the figure. The gray shaded volume is not filled by the liquid, leading to the gravity subtraction (second term in second-to-last line of eq. (3).)

In the latter case, the elastic energy as a function of the Fourier-transformed height profile h_q takes the form:

$$\mathcal{E}[h] = \frac{1}{2} \int \frac{dq}{2\pi} \,\epsilon_q \,|h_q|^2 \,, \qquad \epsilon_q = \gamma \sqrt{q^2 + \kappa^2} \,, \qquad (1)$$

with κ the inverse capillary length, and up to terms of order h^3 . In other cases, interpolation formulas have been proposed [5].

There is numerical evidence [6] that the contact-line energy (1) does not correctly describe the shape of the contact line depinning from a single defect. Surprisingly, the form of ϵ_q as given by (1) seems not to have been questioned, rather the discrepancies were attributed to higherorder terms in h. The latter are indeed present [7,8,9], even for $\theta = 90^{\circ}$ and a vertical wall ($\varphi = 0$), but for small perturbations they are subdominant w.r.t. the dominant term (1).

 $\mathbf{2}$

In the first part of the present article (section 3) we show that for contact-angles leading to a non-flat profile as plotted in figure 1, or for an inclined wall in figure 2, the elastic energy (1) cannot be used, but must be replaced by the more general form

$$\frac{\epsilon_q}{\kappa\gamma} = \frac{\sin(\theta)\cos(\varphi)}{t} + \frac{\left(r^2 - 1\right)\left[t(r+t) + 1\right]\sin^2(\theta)}{t\left(r^2 + 3rt + 3t^2 - 1\right)} \quad (2)$$
$$t = \sqrt{\frac{\sin(\theta + \varphi) + 1}{2}} \quad , \quad r = \sqrt{1 + \frac{q^2}{\kappa^2}} \; .$$

Then we discuss how this form can be measured from the contact line profile in presence of a single defect. This section is pedagogical and self-contained and can be read with no prior knowledge in either wetting or disordered systems.

In the second part of the paper (section 4) we examine the consequences of the form (2) of the elastic kernel ϵ_q for the avalanche statistics of a contact-line at depinning, i.e. a contact-line advancing on a disordered substrate, when the fluid level is adiabatically increased. We calculate the local (i.e. at a given point) as well as global avalanche-size distribution, including the scaling functions. We work at quasi-static depinning using an over-damped equation of motion. The point of the present work is to study the effects arising from the precise form of the elastic kernel. We therefore assume that details of the dynamics as well as non-linearities can be neglected in comparison. We use the methods introduced in [10] to compute the avalanche-size distribution from the Functional Renormalization Group (FRG) theory of pinned elastic systems (see also Refs. [11] for a pedagogical introduction and [12,13] for early numerical tests of this theory). We recall and analyze the formula given in these references, hence this section can be read with no prior knowledge of disordered systems; however understanding its foundations requires the knowledge of the above mentioned literature. Then we derive a general formula for the size distribution for an arbitrary elastic kernel, not given priorly, and compute universal ratios of local-avalanche-size moments versus global ones. Finally, we apply our general result to the contact line with the elasticity (2).

2 Model

Consider a fluid in a semi-infinite reservoir, bounded by a planar plate, in presence of gravity. The plate is inclined by an angle $-\pi/2 < \varphi < \pi/2$ with respect to the vertical, see figure 2. We consider the coordinate system X, y, Z where Z is along the vertical direction, and the equation of the plate is $Z = -X \cot \varphi$. The contact line of the fluid is parameterized by (h(y), y) along the plate, hence

it is at $(X = -h(y) \sin \varphi, y, Z = h(y) \cos \varphi)$ in our coordinate system. The fluid occupies the space Z < Z(X, y) and $X > -Z \tan \varphi$, hence also $X > -h(y) \sin \varphi$, where Z(X,y) is the height of the fluid-air surface. We choose the coordinate Z so that the reservoir level is Z = 0 hence $Z(X \to \infty, y) = 0$.

The energy E is the sum of the fluid-air interface energy, proportional to the area, and the gravitational energy. It is a functional of Z(X, y), with explicit dependence on h(y). Its full expression reads:

$$\begin{aligned} \mathcal{E}[Z,h] &:= \frac{E[Z,h]}{\gamma} \\ &= \int \mathrm{d}y \int_{X>-h(y)\sin\varphi} \mathrm{d}X \bigg[\sqrt{1 + [\partial_X Z]^2 + [\partial_y Z]^2} + \frac{1}{2} \kappa^2 Z^2 \bigg] \\ &- \frac{1}{2} \kappa^2 \int \mathrm{d}y \int_{-h(y)\sin\varphi}^0 \mathrm{d}X \, X^2 \, \cot^2\varphi \\ &- \int \mathrm{d}y \int_{X>0} \mathrm{d}X \, 1 \\ &- \cos\theta \int \mathrm{d}y \, h(y) \;. \end{aligned}$$
(3)

 γ is the surface tension, and

$$\kappa = \frac{1}{L_c} = \sqrt{\frac{\rho g}{\gamma}} \tag{4}$$

defines the capillary length L_c . We call \mathcal{E} the reduced energy. The first term is the area of the fluid surface; the second is the gravitational cost in potential energy for bringing a fluid element from infinity at level Z = 0, filling the reservoir up to height Z(X, y). The second line takes into account that for $\varphi > 0$ and h(y) > 0 (resp. $\varphi < 0$ and h(y) < 0 the volume element $0 < Z < -X \cot \varphi$ (resp. $-X \cot \varphi < Z < 0$ is actually not filled by the liquid; this is the gray shaded region on figure 2. Similarly for $\varphi > 0$ and h(y) < 0 (resp. $\varphi < 0$ and h(y) > 0) there is a corresponding missing volume element which must be added. The third line is the subtraction of the surface energy of the flat profile h(y) = 0 (i.e. Z = 0 and X > 0), necessary to make the problem well-defined. The last term comes from the difference between solid-fluid $\gamma_{\rm SF}$ and solid-air $\gamma_{\rm SA}$ surface energies which defines the *equilibrium* contact angle, denoted θ , via the usual relation $\gamma \cos \theta = \gamma_{\rm SA} - \gamma_{\rm SF}$. The definition of the contact line implies the additional boundary condition

$$Z\left(-h(y)\sin\varphi, y\right) = h(y)\cos\varphi . \tag{5}$$

The energy of a given configuration h(y) of the contact line is obtained as $E[h] := E[Z_h, h]$ where $Z_h(X, y)$ minimizes $\mathcal{E}[Z, h]$ at fixed h(y) under the constraint (5). By translational invariance, the minimum-energy configuration of the contact-line itself, i.e. the minimum of E[h], is attained for a straight line $h(y) = h_0$ where h_0 denotes the equilibrium height.

In the next Section we compute the elastic energy of the contact line, i.e. E[h] to second order in its deformations, i.e. $E[h] - E[h_0] = E_{el}[h] + \mathcal{O}(h^3)$ with

$$E_{\rm el}[h] = \frac{1}{2} \int_q \epsilon_q h_{-q} h_q \ . \tag{6}$$

Here $h(y) = \int_q h_q e^{iqy}$ and we denote $\int_q := \int \frac{dq}{2\pi}$. For a uniform deformation h(y) = h it takes the form:

$$E_{\rm el}[h] = \frac{1}{2}m^2(h - h_0)^2 L_y \quad , \quad m^2 = \epsilon_{q=0} \; , \qquad (7)$$

which defines what we call the mass m, i.e. m^2 is the curvature of the parabolic well in which the contact line sits because of gravity. We then calculate ϵ_q , already announced in eq. (2).

3 Contact-line elasticity

3.1 Model in shifted coordinates

We start by introducing a more convenient expression for the energy of the system. The constraint $X > -h(y) \sin \varphi$ in the domain of integration is tedious to handle, so we introduce the function z(x, y) as

$$z(x,y) := Z(X = x - \hat{h}(y)\sin\varphi, y) . \tag{8}$$

It satisfies the same boundary condition $z(\infty, 0) = 0$. We have also defined

$$\tilde{h}(y) := h(y) - h_0 \tag{9}$$

i.e. z is still the height along the vertical axis, but we have shifted the X coordinate so that the integration domain is

$$x > x_0 := -h_0 \sin \varphi \ . \tag{10}$$

Using that the derivatives, evaluated at $X = x - \tilde{h}(y) \sin \varphi$, satisfy

$$\partial_X Z(X, y) = \partial_x z(x, y) \tag{11}$$

$$\partial_y Z(X,y) = \partial_y z(x,y) + \tilde{h}'(y) \sin \varphi \ \partial_x z(x,y) \ , \qquad (12)$$

one finds that the energy is now a functional noted $E[z, h] = \gamma \mathcal{E}[z, h]$ of z(x, y) and h(y) with

$$\mathcal{E}[z,h] = \int_{y} \mathrm{d}y \int_{x>x_{0}} \mathrm{d}x \left[\frac{\kappa^{2}}{2} z(x,y)^{2} - 1 \right]$$
$$+ \sqrt{1 + [\partial_{x} z(x,y)]^{2} + [\partial_{y} z(x,y) + \tilde{h}'(y) \sin \varphi \, \partial_{x} z(x,y)]^{2}} \right]$$
$$- \frac{\kappa^{2}}{6} \cos^{2} \varphi \, \sin \varphi \int_{y} h(y)^{3}$$
$$+ (\sin \varphi - \cos \theta) \int_{y} h(y) \, . \tag{13}$$

To derive this result, we have used the relation

$$\int_{X>-h(y)\sin\varphi} dX \sqrt{1 + [\partial_X Z]^2 + [\partial_y Z]^2} - \int_{X>0} dX \ 1$$
$$= h(y)\sin\varphi + \int_{X>-h(y)\sin\varphi} dX \sqrt{1 + [\partial_X Z]^2 + [\partial_y Z]^2} - 1 \ ,$$

which was then rewritten in terms of x.

The new height function obeys the constraint

$$z(x_0, y) = h(y)\cos\varphi , \qquad (14)$$

i.e. it is specified on the edge $x = x_0$.

3.2 Zero mode and calculation of the mass

Let us first consider a uniform displacement of the line h(y) = h. We can then restrict to y-independent height functions z(x, y) = z(x), and the reduced energy takes the form

$$L_y^{-1}\mathcal{E}[z,h] = \int_{x>x_0} \mathrm{d}x \left[\sqrt{1+z'(x)^2} - 1 + \frac{\kappa^2}{2} z(x)^2 \right] - \frac{\kappa^2}{6} h^3 \cos^2\varphi \sin\varphi + h(\sin\varphi - \cos\theta) \ . \ (15)$$

There are additional constraints at the boundary,

$$z(x = x_0) = h\cos\varphi , \qquad (16)$$

and z = 0 at infinity. h is arbitrary and not necessarily the preferred value at equilibrium noted h_0 .

The linear variation of the reduced functional $\mathcal{E}[z,h]$ around an arbitrary configuration (z(x),h), by $(\delta z(x), \delta h)$ can be written upon integration by part as

$$\begin{split} \delta \mathcal{E}[z,h]/L_y \\ &= \int_{x > x_0} \mathrm{d}x \, \delta z(x) \bigg[\kappa^2 z(x) - \partial_x \frac{z'(x)}{\sqrt{1 + z'(x)^2}} \bigg] \\ &+ \delta z(x_0) \cos(\theta(x_0) + \varphi) - \frac{\kappa^2}{2} \cos^2 \varphi \sin \varphi \, h^2 \delta h \\ &+ \delta h(\sin \varphi - \cos \theta) \;. \end{split}$$
(17)

We have defined

$$\cos(\theta(x) + \varphi) = -\frac{z'(x)}{\sqrt{1 + z'(x)^2}} \tag{18}$$

$$\sin(\theta(x) + \varphi) = \frac{1}{\sqrt{1 + z'(x)^2}};$$
 (19)

hence $\theta(x_0)$ is the contact angle at the wall and $\theta(x) + \varphi$ is the local angle with respect to the vertical. Note that, because of $\kappa^2 > 0$, the profile decays and the boundary term at infinity does not contribute. The constraint on the boundary implies the additional relation

$$\delta z(x_0) = \delta h \cos \varphi . \tag{20}$$

Consider now the function $z(x) = z_h(x)$ which minimizes the energy at fixed h, i.e. $\delta h = 0$ and $\delta z(x_0) = 0$. This leaves the variations in the bulk, given in the first line of (17), leading to the stationarity condition

$$\kappa^2 z(x) = \frac{z''(x)}{(1+z'(x)^2)^{3/2}} \ . \tag{21}$$

It can be integrated once,

4

$$\frac{\kappa^2}{2}z(x)^2 = 1 - \frac{1}{\sqrt{1 + z'(x)^2}} , \qquad (22)$$

where the integration constant was fixed by considering the limit of $x \to \infty$. This yields at $x = x_0$,

$$\frac{\kappa^2}{2}h^2\cos^2\varphi = 1 - \sin(\theta(x_0) + \varphi)$$
$$= 2\sin^2\left(\frac{\theta(x_0)}{2} + \frac{\varphi}{2} - \pi/4\right) , \quad (23)$$

using that $1 - \sin x = 2\sin^2(x/2 - \pi/4)$. The sign of the root $h \cos \varphi$ must be opposite to the sign of $\theta(x_0) + \varphi - \pi/2$, and similarly when solving for z in eq. (22). This yields

$$h\cos\varphi = -2L_c\sin\left(\frac{1}{2}(\theta(x_0) + \varphi - \frac{\pi}{2})\right) .$$
 (24)

Integrating once more, we obtain the height profile $z = z_h(x)$ in the inverse form,

$$x(z) = x_0 + L_c \left[\operatorname{arcosh} \left(\frac{2L_c}{z} \right) - \operatorname{arcosh} \left(\frac{2L_c}{h \cos \varphi} \right) \right]$$
$$-L_c \left(\sqrt{4 - \frac{z^2}{L_c^2}} - \sqrt{4 - \frac{h^2 \cos^2 \varphi}{L_c^2}} \right) . \tag{25}$$

The integration constant was chosen to satisfy the constraint (16).

The reduced energy of the uniform deformation h(y) = h is thus a simple function $\mathcal{E}(h) = \mathcal{E}[z_h, h]$, whose derivative, $\mathcal{E}'(h)$ is easy to obtain. Indeed we can use the general variational formula (17) around $(z_h(x), h)$ setting $\delta z(x) = \delta h \partial_h z_h(x)$ with $\partial_h z_h(x_0) = 1$ from the constraint (16). The bulk contribution is zero, due to the "equation of motion" (21) satisfied by $z_h(x)$. The boundary contributions, i.e. the second and third lines in (17), give

$$\mathcal{E}'(h)/L_y = \cos\varphi\cos(\theta(x_0) + \varphi) - \frac{\kappa^2}{2}h^2\cos^2\varphi\sin\varphi + \sin\varphi - \cos\theta .$$
(26)

Using eq. (23) this simplifies to

$$\mathcal{E}'(h)/L_y = \cos\theta(x_0) - \cos\theta , \qquad (27)$$

where $\theta(x_0)$ is an implicit function of h, using (24). We first note that this allows to recover the usual condition for equilibrium, $\mathcal{E}'(h_0) = 0$ namely that the local contact



Fig. 3. The height profile z(x), in units of the capillary length L_c , for $\theta = 30^\circ$ and $\varphi = 0$. Different contact-angles $0 < \theta < \pi/2$ are obtained by moving the graph left/right. The case $\pi/2 < \theta < \pi$ is obtained by the reflection $z \to -z$.

angle $\theta(x_0) = \theta$, the equilibrium contact angle. The equation determining h_0 is thus (24), with $\theta(x_0) = \theta$,

$$h_0 \cos \varphi = -2L_c \sin \left(\frac{1}{2}(\theta + \varphi - \frac{\pi}{2})\right)$$
 (28)

Inverting eq. (28) one finds from (27) away from equilibrium

$$\frac{\mathcal{E}'(h)}{L_y} = h \kappa \cos^2 \varphi \sqrt{1 - \frac{1}{4} \kappa^2 h^2 \cos^2 \varphi} + \left(1 - \frac{1}{2} h^2 \kappa^2 \cos^2 \varphi\right) \sin \varphi - \cos \theta .$$
(29)

The mass for the zero mode is thus, after various simplifications

$$m^{2} = \frac{\gamma \mathcal{E}''(h_{0})}{L_{y}} = \frac{\gamma \kappa \sqrt{2} \cos \varphi \sin \theta}{\sqrt{1 + \sin(\theta + \varphi)}} .$$
(30)

An interesting special case is that of a flat interface, $\theta + \varphi = \frac{\pi}{2}$. Then

$$m^2 = \gamma \kappa \sin^2(\theta) . \tag{31}$$

3.3 Elastic energy for arbitrary deformations

We now go back to the general case for h(y). The reduced energy functional $\mathcal{E}[z,h]$ given in (13) explicitly depends on h(y). In addition h(y) also enters in the boundary condition (14). It is important to distinguish these two dependences.

We call $z_h(x, y)$ the profile which minimizes the reduced energy functional $\mathcal{E}[z, h]$ with fixed h(y), hence with respect to bulk variations $\delta z(x, y)$ only,

$$\frac{\delta \mathcal{E}}{\delta z(x,y)|_{\text{bulk}}}[z_h,h] = 0 , \qquad (32)$$

i.e. a partial (functional) derivative w.r.t. z only.

We now expand around the equilibrium solution as follows:

$$h(y) = h_0 + \dot{h}(y)$$
(33)
$$z_h(x, y) = z_0(x) + \tilde{z}(x, y) ,$$

where $z_0(x) := z_{h_0}(x)$ is the equilibrium profile determined in the previous section. Since \tilde{z} is of order $\mathcal{O}(\tilde{h})$, to compute the elastic energy we need to expand the minimum energy to second order in \tilde{h} and \tilde{z} . Expanding (13) to second order in the *explicit* dependence on \tilde{z} and \tilde{h} we have

$$\mathcal{E}[z_h, h] - \mathcal{E}[z_0, h_0] = \delta^{(1)} \mathcal{E} + \delta^{(2)} \mathcal{E} + \mathcal{O}(\tilde{h}^3) \quad (34)$$

$$\delta^{(2)} \mathcal{E} = \delta^{(2,1)} \mathcal{E} + \delta^{(2,2)} \mathcal{E} + \delta^{(2,3)} \mathcal{E} .$$

The first term is the linear variation,

$$\delta^{(1)}\mathcal{E} = \frac{\delta E}{\delta z}[z_0, h_0] \cdot \tilde{z} + \frac{\delta E}{\delta h}[z_0, h_0] \cdot \tilde{h} , \qquad (35)$$

whereas $\delta^{(2,p)} \mathcal{E}$ are second order variations computed below. Let us start with $\delta^{(1)} \mathcal{E}$:

$$\delta^{(1)}\mathcal{E} = \int_{y} \int_{x > x_{0}} \tilde{z}(x, y) \left[\kappa^{2} z_{0}(x) - \partial_{x} \frac{z_{0}'(x)}{\sqrt{1 + z_{0}'(x)^{2}}} \right] + \int_{y} \delta z(x_{0}, y) \cos(\theta(x_{0}) + \varphi) - \frac{\kappa^{2}}{2} \cos^{2} \varphi \sin \varphi h_{0}^{2} \int_{y} \tilde{h}(y) + \int_{y} \tilde{h}(y) (\sin \varphi - \cos \theta) .$$
(36)

The first line is the bulk variation which vanishes since $z_0(x)$ satisfies the stationarity condition (21). The rest is the sum of the *boundary* variation in z and the variation in h. Thanks to the exact constraint

$$\tilde{z}(x_0, y) = \tilde{h}(y) \cos \varphi , \qquad (37)$$

one can check, using the results of the previous section, that this sum vanishes identically if h_0 is the equilibrium value. Hence we have $\delta^{(1)}\mathcal{E} = 0$. Since \tilde{z} also contains subdominant \tilde{h}^2 contributions, this is a quite useful observation, as we do not need to worry about them in computing the elastic energy. They drop out by virtue of the equilibrium condition.

Let us now study the three second variations. The first one is

$$\delta^{(2,1)}\mathcal{E} = \frac{1}{2}\tilde{z} \cdot \frac{\delta^2 E}{\delta z \delta z} [z_0, 0] \cdot \tilde{z}$$
(38)
$$= \frac{1}{2} \int_{y,x>x_0} \kappa^2 \tilde{z}(x,y)^2 + \frac{[\partial_x \tilde{z}(x,y)]^2}{[1+z_0'(x)^2]^{3/2}} + \frac{[\partial_y \tilde{z}(x,y)]^2}{[1+z_0'(x)^2]^{1/2}} .$$

The second one is

δ

$$^{(2,2)}\mathcal{E} = \tilde{h} \cdot \frac{\delta^2 E}{\delta h \delta z} [z_0, h_0] \cdot \tilde{z}$$

$$= \sin \varphi \int_{y, x > x_0} h'(y) \frac{z'_0(x)}{\sqrt{1 + z'_0(x)^2}} \partial_y \tilde{z}(x, y) ,$$
(39)

where we have used $\tilde{h}'(y) = h'(y)$ to alleviate the notation. The third contribution is

$$\delta^{(2,3)}\mathcal{E} = \frac{1}{2}\tilde{h} \cdot \frac{\delta^2 E}{\delta h \delta h} [z_0, h_0] \cdot \tilde{h}$$
$$= \delta^{(2,3a)}\mathcal{E} + \delta^{(2,3b)}\mathcal{E} , \qquad (40)$$

where we write separately the second derivative with respect to the explicit dependence on $\tilde{h}(y)$ of the surface energy, namely

$$\delta^{(2,3a)}\mathcal{E} = \frac{1}{2}\sin^2\varphi \int_y \int_{x>x_0} h'(y)^2 \frac{z'_0(x)^2}{\sqrt{1+z'_0(x)^2}} ,\quad (41)$$

and of the gravitational one,

$$\delta^{(2,3b)}\mathcal{E} = -\frac{1}{2}\kappa^2 \cos^2 \varphi \,\sin\varphi \,h_0 \int_y \tilde{h}(y)^2 \,. \tag{42}$$

To compute the first two contributions one needs to specify the properties of $\tilde{z}(x, y)$ which follow from its definition (33). The function $z_h(x, y)$ must obey equation (32), for any h. Hence we can expand this equation order by order in h. For h = 0 it yields again the stationarity condition (21) for $z_0(x)$; to first order it yields

$$\tilde{z} \cdot \frac{\delta \mathcal{E}}{\delta z \delta z(x,y)|_{\text{bulk}}} [z_0, h_0] + \tilde{h} \cdot \frac{\delta \mathcal{E}}{\delta h \delta z(x,y)|_{\text{bulk}}} [z_0, h_0] = 0$$

which, in explicit form, yields the equation obeyed by $\tilde{z}(x, y)$ (to linear order in h which is all we need here):

$$\begin{bmatrix} \kappa^2 - \partial_x \frac{1}{[1 + z_0'(x)^2]^{3/2}} \partial_x - \frac{1}{[1 + z_0'(x)^2]^{1/2}} \partial_y^2 \end{bmatrix} \tilde{z}(x, y) \\ = \sin\varphi \, h''(y) \frac{z_0'(x)}{[1 + z_0'(x)^2]^{1/2}} \,. \tag{43}$$

It must be solved with the boundary condition (37) and $\tilde{z}(x = \infty, y) = 0$.

Remarkably, this complicated looking equation, which depends on the profile $z_0(x)$ known only in the implicit form (25), can be solved analytically. First of all, using the stationarity equation (21) satisfied by $z_0(x)$, one notes that

$$\tilde{z}(x,y) = \tilde{z}_1(x,y) := -\sin\varphi \ h(y)z'_0(x)$$
 (44)

is a particular solution of (43) which vanishes at $x = \infty$ and takes the value

$$\tilde{z}_1(x_0, y) = \sin\varphi \cot(\theta + \varphi)h(y) \tag{45}$$

at the boundary. The full solution can thus be written as

$$\tilde{z}(x,y) = \tilde{z}_1(x,y) + \tilde{z}_2(x,y) ,$$
 (46)

where $\tilde{z}_2(x, y)$ satisfies the homogeneous equation, i.e. (43) setting the r.h.s. to zero. The boundary condition (14) implies

$$\tilde{z}_2(x_0, y) = \frac{\sin\theta}{\sin(\theta + \varphi)} h(y) .$$
(47)

Both \tilde{z}_1 and \tilde{z}_2 vanish at $x = \infty$.

To solve the homogeneous equation, we go to Fourier space in y-direction, and introduce a new variable in x-direction. We thus look for the solution in the form

$$\tilde{z}_2(x,y) = \frac{\sin\theta}{\sin(\theta+\varphi)} \int_q e^{iqy} \tilde{h}(q) F_{\tilde{q}}(S(x))$$
(48)

$$S(x) = \sin(\theta(x) + \varphi) = \frac{1}{\sqrt{1 + z'_0(x)^2}}$$
(49)

$$\tilde{q} = q/\kappa \tag{50}$$

with $S(x_0) = \sin(\theta + \varphi)$. Deriving eq. (49) w.r.t. x, and using (21) and (22) to express the result in terms of S(x), we obtain the rule for changing the derivatives,

$$\partial_x = \kappa \frac{\sqrt{2}(1-S)\sqrt{S+1}}{S} \partial_S . \tag{51}$$

The resulting equation for the function $F_{\tilde{q}}(S)$ reads

$$\begin{aligned} (\tilde{q}^2 S + 1) F_{\tilde{q}}(S) + (1 - S)(7S^2 + S - 4) F'_{\tilde{q}}(S) \\ -2(S - 1)^2 S(S + 1) F''_{\tilde{q}}(S) &= 0 , \end{aligned}$$
(52)

where $\sin(\theta + \varphi) < S < 1$. The constraint (47) implies the boundary conditions $F_{\tilde{q}}(\sin(\theta + \varphi)) = 1$ and $F_{\tilde{q}}(1) = 0$. After some search, the general solution of eq. (52) which satisfies $F_{\tilde{q}}(1) = 0$ is found to be

$$F_{\bar{q}}(S) = \frac{g_r(S)}{g_r(\sin(\theta + \varphi))}$$
(53)
$$g_r(S) = \frac{(1 - S)^{r/2}}{S} \left(1 + \frac{\sqrt{S+1}}{\sqrt{2}}\right)^{-r} \times \left(2r^2 + 3\sqrt{2}r\sqrt{S+1} + 3S + 1\right) ,$$
(54)

where we denoted

 $\mathbf{6}$

$$r := \sqrt{1 + \tilde{q}^2} \ . \tag{55}$$

We now evaluate the second variation. Consider first the sum of (38) and (39). Using the equation of motion (43) for \tilde{z} , the combination $\delta^{(2,1)}\mathcal{E} + \frac{1}{2}\delta^{(2,2)}\mathcal{E}$ can be integrated by part. Therefore, we obtain for the combination δ $[\delta^{(2,1)}\mathcal{E} + \frac{1}{2}\delta^{(2,2)}\mathcal{E}] + \frac{1}{2}\delta^{(2,2)}\mathcal{E}$

$$\delta^{(2,1)}\mathcal{E} + \delta^{(2,2)}\mathcal{E} = -\frac{1}{2} \int_{y} \frac{\tilde{z}(x_{0}, y)}{[1 + z_{0}'(x)^{2}]^{3/2}} \partial_{x} \tilde{z}(x, y) \Big|_{x=x_{0}} -\frac{1}{2} \sin \varphi \int_{y, x > x_{0}} h''(y) \frac{z_{0}'(x)}{\sqrt{1 + z_{0}'(x)^{2}}} \tilde{z}(x, y) .$$
(56)

The last term has been integrated by part w.r.t. y^{2} . To continue, we note the useful equality

$$-\int_{y} \frac{\tilde{z}_{1}(x_{0}, y)}{[1 + z_{0}'(x)^{2}]^{3/2}} \partial_{x} \tilde{z}_{2}(x, y)\Big|_{x=x_{0}}$$

$$= -\int_{y} \frac{\tilde{z}_{2}(x_{0}, y)}{[1 + z_{0}'(x)^{2}]^{3/2}} \partial_{x} \tilde{z}_{1}(x, y)\Big|_{x=x_{0}}$$

$$+ \sin \varphi \int_{y, x > x_{0}} h''(y) \frac{z_{0}'(x)}{\sqrt{1 + z_{0}'(x)^{2}}} \tilde{z}_{2}(x, y) , \quad (57)$$

which is a consequence of the two different ways to integrate by part

$$\begin{split} \int_{y,x>x_0} \left[\kappa^2 \tilde{z}_1(x,y) \tilde{z}_2(x,y) + \frac{[\partial_x \tilde{z}_2(x,y)] [\partial_x \tilde{z}_1(x,y)]}{[1+z_0'(x)^2]^{3/2}} \right. \\ \left. + \frac{[\partial_y \tilde{z}_1(x,y)] [\partial_y \tilde{z}_2(x,y)]}{[1+z_0'(x)^2]^{1/2}} \right] \,, \end{split}$$

² Note that the integration by part of ∂_y produces no surface term. This can be made rigorous considering a periodic modulation h(y). We thus restrict here to functions that can be written as sum of periodic modulations, or have compact support.

and to use the equation of motion for \tilde{z}_1 (inhomogeneous) and \tilde{z}_2 (homogeneous).

Inserting $\tilde{z} = \tilde{z}_1 + \tilde{z}_2$ into (56) and using the equality (57) we get

$$\delta^{(2,1)}\mathcal{E} + \delta^{(2,2)}\mathcal{E} = -\frac{1}{2} \int_{y} \frac{\tilde{z}_{2}(x_{0}, y)}{[1 + z_{0}'(x)^{2}]^{3/2}} \partial_{x} \tilde{z}_{2}(x, y) \Big|_{x=x_{0}} -\frac{1}{2} \int_{y} \frac{\tilde{z}_{1}(x_{0}, y)}{[1 + z_{0}'(x)^{2}]^{3/2}} \partial_{x} \tilde{z}_{1}(x, y) \Big|_{x=x_{0}} -\int_{y} \frac{\tilde{z}_{2}(x_{0}, y)}{[1 + z_{0}'(x)^{2}]^{3/2}} \partial_{x} \tilde{z}_{1}(x, y) \Big|_{x=x_{0}} -\frac{1}{2} \sin \varphi \int_{y, x > x_{0}} h''(y) \frac{z_{0}'(x)}{\sqrt{1 + z_{0}'(x)^{2}}} \tilde{z}_{1}(x, y) .$$
(58)

We now discuss simplifications. Firstly, the last term in eq. (58) exactly cancels $\delta \mathcal{E}^{(2,3b)}$; this is shown using (44). Secondly, from (44), (21) and (14), we obtain

$$\frac{\partial_x \tilde{z}_1(x,y)}{[1+z_0'(x)^2]^{3/2}}\Big|_{x=x_0} = -\sin\varphi\cos\varphi\kappa^2 h(y)h_0 \ . \tag{59}$$

This shows that the second line, half the third line and $\delta^{(2,3b)}\mathcal{E}$ cancel; the remaining half of the third line gives the first term reported in eq. (60) below. The second term comes from the first line of (58), using (48), so we get finally:

$$5^{(2)}\mathcal{E} = \frac{\sin\varphi\cos\varphi\sin\theta}{2\sin(\theta+\varphi)}\kappa^2 h_0 \int_y h(y)^2$$
(60)
$$-\frac{1}{2}\sin^2\theta\sin(\theta+\varphi) \int_q h_q h_{-q} F_{\tilde{q}}(S(x))\partial_x F_{\tilde{q}}(S(x))\Big|_{x=x_0} .$$

To compute the second term we use rule (51), where at the end S must be evaluated on the boundary $S = \sin(\theta + \varphi)$. To compute the first term we use the value (23) for h_0 .

This yields our final result for the elastic energy,

$$E_{\rm el}[h] = \frac{1}{2} \int_q \epsilon_q h_q h_{-q} , \qquad (61)$$

with

$$\frac{\epsilon_q}{\kappa\gamma} = \frac{\sin(\theta)\cos(\varphi)}{t} + \frac{(r^2 - 1)[t(r+t) + 1]\sin^2(\theta)}{t(r^2 + 3rt + 3t^2 - 1)}$$
$$t = \sqrt{\frac{\sin(\theta + \varphi) + 1}{2}} \quad , \quad r = \sqrt{1 + \frac{q^2}{\kappa^2}} \quad . \tag{62}$$

One finds that ϵ_q is a scaling function of q/κ which reproduces formula (30) for the energy of a uniform mode $\epsilon_{q=0} = m^2$ as computed in the previous section, and which behaves as $\epsilon_q \approx \kappa \gamma \sin^2 \theta |q|$ for large |q|.

When $\varphi + \theta = \pi/2$, the equilibrium shape of the interface is flat. Thus the elastic energy is expected to simplify. Indeed, it becomes

$$\frac{\epsilon_q}{\kappa\gamma} = \sin^2(\theta)\sqrt{1 + \frac{q^2}{\kappa^2}} . \tag{63}$$



Fig. 4. $\kappa\gamma/\epsilon_q$ as a function of qL_c , for $\theta = 40^\circ$, $\varphi = 0^\circ$, $10^\circ, \ldots, 90^\circ$ (from bottom to top).



Fig. 5. Sample profiles h(y), in units of $f/(\kappa\gamma)$, plotted as a function of y/L_c , for $\theta = 40^\circ$, $\varphi = 0^\circ$ (bottom) and $\varphi = 45^\circ$ (top). The profiles are the response to a force inside the green box $|y| < \delta = 0.3L_c$. The profile has to be continued symmetrically to the left.

3.4 The contact-line profile as a means of measuring ϵ_q

Suppose the contact line is in force-free equilibrium. Then pull on it with a force per unit length:

$$F(y) = \frac{f}{2\delta}\theta(|y| < \delta) .$$
 (64)

This leads in linear response (i.e. for the quadratic in h elastic energy we are using) to the following profile

$$h(y) = f \int_0^\infty \frac{dq}{\pi} \frac{\cos(qy)}{\epsilon_q} \frac{\sin(q\delta)}{q\delta} .$$
 (65)

The limit of a δ -like force is recovered in the limit of $\delta \to 0$, which eliminates the last factor of $\sin(q\delta)/(q\delta)$. However the latter makes the integral convergent at large q. We have plotted for $\theta = 40^{\circ}$ and $\delta = 0.3$ two solutions on figure 5, one for $\varphi = 0^{\circ}$ (bottom) and the other for $\varphi = 45^{\circ}$ (top). We note that ϵ_q can be reconstructed from the profile as follows

$$\epsilon_q = f \frac{\sin(q\delta)}{q\delta} \left[2 \int_0^\infty \mathrm{d}y \cos(qy) h(y) \right]^{-1} .$$
 (66)



Fig. 6. Top: Blue points: Reconstruction of ϵ_q , in units of $\kappa\gamma$, as a function of q/κ , using (66) from the points of the bottom curve of figure 5, i.e. a vertical plate geometry $\varphi = 0$. Solid red line: the analytical result. One sees a numerical problem appearing at large q, at $q = \pi/\delta$, where δ is the box-size in figure 5. However this is already far in the linear asymptotic regime (dashed line). Bottom: The same plot for smaller q. One sees that ϵ_q is well reconstructed.

A reconstruction of ϵ_q starting from the blue points on the top of figure 5 is given in figure 6.

4 Avalanche-size distributions

4.1 The model

We now study the case of a disordered plate, which is immersed with velocity v into the liquid reservoir. This is the geometry of the experiment described in Ref. [1]. We use some of the notations defined there: x denotes the coordinate along the contact line (denoted y in the previous section) and u(x) the height of the contact line along the plate, in the frame of the plate. We reserve the notation h(x) to the contact-line height measured in the laboratory frame, with the relation³:

$$h(x,t) = u(x,t) - w$$
, $w = vt$. (67)

 $^{^{3}\,}$ In the experiment of Ref. [1], there is an additional slowly varying offset.

Even though it is a dynamical problem, it is useful to introduce the energy

$$\mathcal{H}[u] = \int_0^L \mathrm{d}x \, \frac{m^2}{2} \left[u(x) - w \right]^2 + V(x, u(x)) + \delta E[u] \,. \tag{68}$$

Here m^2 is the mass of the zero mode q = 0, and $\delta E[u] := \frac{1}{2} \int_q [\epsilon_q - \epsilon_0] u_q u_{-q}$ the remaining part of the elastic energy at non-zero wave vector $q \neq 0$. For the contact-line, $m^2 = \epsilon_0$ and $\delta E[u]$ are both given in (62). We neglect possible non-linear elastic terms [8,9]. The function V(x, u) is a random potential, whose derivative $\partial_u V(x, u)$ is short-ranged correlated, modeling the disordered substrate.

The contact line is pinned by the disorder, but also trapped in the quadratic well with curvature m^2 . Advancing the well-position w by immersing the plate with velocity v leads to a motion of the contact line, which we now study in the quasi-static limit, i.e. the limit of small v. From the energy form (68) one can derive, upon various assumptions about the fluid, equations of motion (see e.g. [4,7] and references therein). Our assumption here is that in the quasi-static limit, they lead to the same statistics as the simplest over-damped model studied in Ref. [10], at least to the level of approximation that we use here (i.e. lowest order, i.e. one-loop FRG). Furthermore, although quasi-static dynamics and pure statics are different, they lead, in the same order of approximation, to identical rescaled avalanche-size distributions. Deviations are expected only at the next, i.e. two-loop order. This justifies our studying of the energy form (68), and, as we will see, our point is that noticeable effects already arise from the precise form of the elastic kernel.

4.2 Global statistics of avalanches

4.2.1 Definitions

As can be seen on figure 8, the experiment [1] shows, as predicted by the theory [12, 10], that the motion of the



Fig. 7. Sketch of the experimental setup used in [1]. The size of the image in the inset is 1.5 mm.



Fig. 8. Data from [1]: Height of the contact line $\bar{h}(w)$ averaged over $2L_c$, as a function of the position w of the plate (system: iso/Si). The fast depinning events (upwards) are clearly visible. Between them, the contact line moves downwards at the plate velocity v (here 1 μ m/s). The straight line is the reference level h_0 .

contact-line proceeds by sudden jumps, i.e. avalanche motion. For an avalanche occurring at a given position w of the center of the quadratic well, one defines $u_w^-(x)$ the contact-line position just before the avalanche and $u_w^+(x)$ its position just after. The size of the avalanche is defined as

$$S := \int_0^L \mathrm{d}x \, \left[u_w^+(x) - u_w^-(x) \right] = \int_0^L \mathrm{d}x \, \left[h_w^+(x) - h_w^-(x) \right] \, .$$
(69)

Hence one can measure the avalanche size S as the area swiped both in the u or in the h coordinate system.

We now recall the main results of Refs. [12,10] and apply the general formulation to various cases, including the contact line for which we have computed the elasticity in section 3. The characteristic rescaled function $\tilde{Z}(\lambda)$ is defined as:

$$\tilde{Z}(\lambda) := \frac{S_m}{\langle S \rangle} \left\langle e^{\lambda S/S_m} - 1 \right\rangle \tag{70}$$

$$S_m := \frac{\langle S^2 \rangle}{2 \langle S \rangle} = \frac{-\Delta'(0^+)}{m^4} . \tag{71}$$

All averages $\langle \ldots \rangle$ are over the normalized probability density P(S) of avalanches. By definition $\tilde{Z}(\lambda) = \lambda + \lambda^2 + \mathcal{O}(\lambda^3)$. The function $\Delta(w)$ is the renormalized correlator of the disorder defined and studied in [12,10] and measured in Ref. [1]. It is *m* dependent: at large *m* it is equal to the bare disorder correlator, while as *m* is decreased it develops a linear cusp at u = 0, whose value, $\Delta'(0^+)$, is related to the second moment of the size distribution as displayed above in equation (71).

The scale S_m is the large-scale cutoff for avalanche sizes, originating from the quadratic well which suppresses the largest avalanches. It is an important scale as it allows to define universal functions in the limit where it becomes large i.e. $S_m \gg S_{\min}$, where S_{\min} is the typical size of the smallest avalanches. In the variable $s := S/S_m$ the avalanche-size distribution becomes universal. Universal means independent of small-scale details, but not of the large-scale setting, e.g. it will depend on the precise form of the elastic kernel. Indeed, one of the predictions of the FRG theory is that if the exponent τ satisfies $2 > \tau > 1$ which will turn out to be the case here, then the distribution of avalanche sizes for $S \gg S_{\min}$ takes the form as $m \to 0$, i.e. $S_m \gg S_{\min}$,

$$P(S)dS := \frac{\langle S \rangle}{S_m} p\left(\frac{S}{S_m}\right) \frac{dS}{S_m} .$$
 (72)

The function p(s) is universal in the above sense. For a fixed elastic kernel it depends only on the space dimension d. Note that the normalized probability P(S) depends on the cut-off S_{\min} via the first moment $\langle S \rangle$ which cannot be predicted by the theory, hence is an input from experiment. It is important to stress that while the function p(s) is universal and convenient for data analysis, *it is not a probability distribution* and is not normalized to unity. Rather, it satisfies from its definition (72) and using (71) the two normalization conditions

$$\langle s \rangle_p = \int \mathrm{d}s \, s \, p(s) = 1 \;, \tag{73}$$

$$\left\langle s^2 \right\rangle_p = \int \mathrm{d}s \, s^2 p(s) = 2 \;.$$
 (74)

Here and below we use the notation $\langle s \rangle_p$ to denote an integration over p(s) and distinguish it from a true expectation value over P(S), denoted $\langle \ldots \rangle$. Note that $\tilde{Z}(\lambda)$ and p(s) are related by the Laplace transform

$$\tilde{Z}(\lambda) = \int_0^\infty \mathrm{d}s \, p(s) \left[e^{\lambda s} - 1 \right] \,. \tag{75}$$

Note finally that the limit $m \to 0$ means that L_c is large compared to the microscopic cutoff *a* along the line, which may be of nanometer scale, or for strong disorder the size of the defects.

4.2.2 Results from one-loop FRG

We now summarize the results obtained in [10] for a general form of the elastic kernel ϵ_k . One defines:

$$\tilde{\epsilon}_k := \epsilon_k / \epsilon_0 , \qquad \epsilon_0 = m^2 .$$
(76)

The general result of [10] is that $Z(\lambda)$ satisfies, in an expansion in powers of the renormalized disorder $\Delta(w)$ and up to terms of $\mathcal{O}(\Delta^2)$,

$$\tilde{Z} = \lambda + \tilde{Z}^2 + \alpha J(\tilde{Z}) , \qquad \alpha = -\epsilon \int_k \epsilon_k^{-2} \Delta''(0^+)$$

$$T(z) = \frac{1}{2} \int_k \left[\frac{z^2}{z^2} - \frac{z}{z^2} - \frac{z^2}{z^2} \right]$$

$$J(z) := \frac{1}{\epsilon \int_k \tilde{\epsilon}_k^{-2}} \int_k \left[\frac{1}{(\tilde{\epsilon}_k - 2z)^2} + \frac{1}{\tilde{\epsilon}_k - 2z} - \frac{1}{\tilde{\epsilon}_k} - 3\frac{1}{\tilde{\epsilon}_k^2} \right].$$
(78)

Note that the parameter

$$\epsilon := d_c - d , \qquad (79)$$

9

the distance to the upper critical dimension, introduced here for later purpose, cancels and is thus immaterial in this formula. As such, this equation is formally exact for any m, any ϵ_k and any dimension, up to $\mathcal{O}(\Delta^2)$ terms.

Furthermore, consider now an elastic kernel such that:

$$\epsilon_k \sim_{k \to \infty} K k^{\beta}$$
 (80)

with elasticity range $\beta \leq 2$ (i.e. long-ranged for $\beta < 2$ and short-ranged for $\beta = 2$) and K the elastic constant. Assume that it takes the form

$$\tilde{\epsilon}_k = e(K^{1/\beta}k/\mu) , \qquad \mu^\beta := m^2 , \qquad (81)$$

where the elastic scaling function e(p) is constructed as the unique dimensionless function of the dimensionless argument p which satisfies e(0) = 1, and $e(p) = p^{\beta}$ for $p \to \infty$. Then, the result of [10] is that the formula (77) can be controlled, in the limit $m \to 0$ as an expansion in powers of $\epsilon = d_c - d$, where $d_c = 2\beta$ is the upper critical dimension. For this one uses that

$$\int_{k} \epsilon_{k}^{-2} = \mu^{d-2\beta} K^{-d/\beta} \int_{p} e(p)^{-2}$$
$$= \mu^{-\epsilon} K^{-2+\frac{\epsilon}{\beta}} \frac{C_{\beta}}{\epsilon} + \mathcal{O}(1)$$
(82)

in the limit $\epsilon = d_c - d \to 0$, where $C_{\beta} = 2(4\pi)^{-d_c/2}/\Gamma(\beta)$ is *independent* of the shape of the function e(p). Since the 1-loop FRG flow for Δ has the schematic form $-\mu\partial_{\mu}\Delta = -\mu\partial_{\mu}(\int_k \epsilon_k^{-2}) \times \Delta^2 = (\epsilon \int_k \epsilon_k^{-2})\Delta^2$, it is convenient to introduce the rescaled disorder,

$$\tilde{\Delta}''(0^+) := \left(\epsilon \int_k \epsilon_k^{-2} \right) \Delta''(0^+) = K^{-2} C_\beta \mu^{-\epsilon} \Delta''(0^+) , \quad (83)$$

where the second equality holds to leading order in ϵ . This is precisely the parameter α defined above, i.e. $\alpha = -\tilde{\Delta}''(0)$. The one-loop FRG flow for the rescaled disorder admits a fixed point, from which one obtains, as $m \to 0$, that α flows to $\alpha = -\tilde{\Delta}''(0) = -\frac{1}{3}\epsilon(1-\zeta_1)$. Here ζ_1 is the $\mathcal{O}(\epsilon)$ correction to the roughness exponent of the elastic object. For the type of disorder relevant for the contact line (i.e. random-field disorder) one has $\zeta_1 = 1/3$, hence

$$\alpha = -\frac{2}{9}\epsilon . \tag{84}$$

This value will be used from now on in (77), which is thus valid up to terms of order $\mathcal{O}(\epsilon^2)$. Using the above definitions one finds that to lowest order in ϵ , J(z) depends only on the dimensionless scaling function e(p), and not on the stiffness K:

$$J(z) = C_{\beta}^{-1} \int_{p} \left[\frac{z^2}{(e(p) - 2z)^2} + \frac{z}{e(p) - 2z} - \frac{z}{e(p)} - 3\frac{z^2}{e(p)^2} \right],$$
(85)

valid for SR elasticity as well as LR elasticity. Of course, since we work to lowest order in ϵ , the above integral should be computed in the critical dimension $d = d_c$.

Note that the mean-field singularity of this equation corresponds to p = 0, e(p = 0) = 1, i.e. z = 1/2.

4.3 From $\tilde{Z}(\lambda)$ **to** p(s)

In Refs. [12, 10] we examined various choices for the elastic kernel ϵ_k , computed $\tilde{Z}(\lambda)$ for each choice, and extracted the scaled avalanche distribution p(s). Here we show that p(s) can directly be written as a function of ϵ_k , or e(p).

4.3.1 General formalism

The definition (75) implies

$$\int_0^\infty ds \, sp(s)e^{\lambda s} = \tilde{Z}'(\lambda) \,. \tag{86}$$

While p(s) does not admit a Laplace transform, the function sp(s) does. Laplace inversion then yields

$$sp(s) = \frac{1}{2i\pi} \int_{-i\infty}^{i\infty} d\lambda \, e^{-\lambda s} \tilde{Z}'(\lambda)$$
$$= \frac{1}{2i\pi} \int_{-i\infty}^{i\infty} dZ \, e^{-s(Z-Z^2 - \alpha J(Z))} , \qquad (87)$$

up to terms of order α^2 . At this stage there is a heuristic step, to go from the λ to the Z contour. We assume that the contour Z = ix with $x \in \mathbb{R}$ is the correct one, as it is for the mean-field case ($\alpha = 0$) around which we perturb. We will check this result on known cases below. Then one has

$$sp(s) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \, e^{-isx - sx^2 + s\alpha J(ix)}$$

$$= sp_{\rm MF}(s) + \frac{\alpha s}{2\pi} \int_{-\infty}^{\infty} dx \, J(ix) e^{-isx - sx^2}$$
(88)

to lowest order in α . We have introduced the scaled meanfield avalanche-size distribution, i.e. (87) at $\alpha = 0$,

$$p_{\rm MF}(s) = \frac{1}{2\sqrt{\pi}} s^{-3/2} e^{-s/4} .$$
 (89)

For the elastic manifold, it holds for $d \ge d_c$. We want to compute the correction to sp(s) of order α , i.e. of order $\epsilon = d_c - d$:

$$\frac{s}{2\pi} \int_{-\infty}^{\infty} dx J(ix) e^{-isx-sx^2}$$

$$= \frac{1}{\epsilon \int_k \tilde{\epsilon}_k^{-2}} \frac{s}{2\pi} \int_0^{\infty} dt \int_k \int_{-\infty}^{\infty} dx \left[t e^{-t(\tilde{\epsilon}_k - 2ix)} \partial_y^2 + e^{-t(\tilde{\epsilon}_k - 2ix)} \partial_y - e^{-t\tilde{\epsilon}_k} \partial_y - 3t e^{-t\tilde{\epsilon}_k} \partial_y^2 \right] e^{-isx-sx^2 + iyx} |_{y=0}$$

$$= \frac{1}{\epsilon \int_k \tilde{\epsilon}_k^{-2}} \frac{s}{2\pi} \int_0^{\infty} dt \int_k \int_{-\infty}^{\infty} dx \, x e^{-\tilde{\epsilon}_k t - isx-sx^2} \times \left[e^{2itx} (i - tx) + 3tx - i \right].$$
(90)

The integrand behaves (before integration over x and k) as t^2 at small t as a result of the counterterms. It is useful to introduce the *elastic generating function*,

$$\mathcal{C}(t) := \frac{1}{\epsilon \int_k \tilde{\epsilon}_k^{-2}} \int_k e^{-t\tilde{\epsilon}_k} = \frac{1}{C_\beta} \int_p e^{-te(p)} , \qquad (91)$$

in terms of which we get, integrating (90) over x:

$$\frac{p(s)}{p_{\rm MF}(s)} = 1 + \alpha \frac{1}{4s} \int_0^\infty dt \ \mathcal{C}(t) X(t,s) \tag{92}$$
$$X(t,s) = e^{t-t^2/s} \Big[4t^3 + s^2(2+t) - 2st(3+2t) \Big]$$
$$-s \Big[2s + 3t(s-2) \Big] \ . \tag{93}$$

This is our final and most general formula, valid for any elasticity ϵ_k and to first order in α , i.e. in ϵ . By performing the integral over s, one checks that it satisfies automatically the two normalization conditions for p(s) (to order α).

4.3.2 Special cases

We now check that this formula recovers previous known results.

Standard local elasticity: For $\epsilon_k = Kk^2 + m^2$ one has $\beta = 2$ and $e(p) = p^2 + 1$. Using that $\int_k e^{-tk^2} = (4\pi)^{-d/2}t^{-d/2}$ and computing the momentum integral in $d = d_c = 4$, one finds

$$C(t) = \frac{1}{2t^2} e^{-t} . (94)$$

Inserting into (92) and performing the t integral yields

$$\frac{p(s)}{p_{\rm MF}(s)} = 1 + \frac{\alpha}{16} \left[(\ln s + \gamma_E)(s-6) + 4s - 8\sqrt{\pi}\sqrt{s} + 4 \right],\tag{95}$$

which recovers the result (169) of [10] (to lowest order in α).

Elasticity of flat contact line and generalization: From (63) the elasticity of a flat contact line $\varphi + \theta = \pi/2$ is $\epsilon_k = \gamma \sin^2 \theta \sqrt{k^2 + \kappa^2}$. This gives:

$$\beta = 1 \quad , \quad K = \gamma \sin^2 \theta \quad , \quad m^2 = \mu = K/\kappa \tag{96}$$

$$e(p) = \sqrt{p^2 + 1}$$
 (97)

It is instructive to slightly generalize this, and study

$$e(p) = \sqrt{p^2 + (1-a)^2} + a \tag{98}$$

with 0 < a < 1, which interpolates between the flat contact line for a = 0 and e(p) = |p| + 1 for a = 1, two cases studied in Ref. [10]. From the definition (91), computing the integral in $d = d_c = 2$, we obtain

$$C(t) = \frac{e^{-t}}{t^2} \left[1 + (1-a)t \right] .$$
(99)

Inserting into (92) and performing the t integral yields

$$\frac{p(s)}{p_{\rm MF}(s)} = 1 + \frac{\alpha}{8} \left\{ 16 - 12a - 6\gamma_{\rm E} + (1-a)\sqrt{\pi}s^{3/2} + \left[(3-2a)s - 6 \right] \ln s - 4(3-a)\sqrt{\pi}\sqrt{s} + s(2a(5-\gamma_{\rm E}) + 3(\gamma_{\rm E} - 2)) \right\}.$$
 (100)

For a = 0, this is the same as (E14) of [10], and for a = 1 has the same form as (95) upon a rescaling of α by a factor of 2, as noted in Ref. [10].

To obtain nicer forms for p(s), and more convenient for extrapolations in physical dimension, one needs to reexponentiate these direct ϵ expansion results, as done in Ref. [10]. We will not attempt to do this here for the general case.

4.4 More on the generating function $\tilde{Z}(\lambda)$

 $Z(\lambda)$ is easier to compare with numerical and experimental data than the disorder distribution p(s), since even if their statistics is mediocre averages over all data are taken, thus the statistical fluctuations are less pronounced. Therefore, we come back to the general formula and some examples.

4.4.1 Generating function in the general case

Since we work to the first order in α , i.e. in $\epsilon = d_c - d$ we can write:

$$\tilde{Z}(\lambda) = \frac{1}{2} \left(1 - \sqrt{1 - 4\lambda} \right) + \alpha \frac{J\left(\frac{1}{2} [1 - \sqrt{1 - 4\lambda}]\right)}{\sqrt{1 - 4\lambda}} \Big|_{z = Z_{\mathrm{MF}(\lambda)}}.$$
(101)

up to higher order terms, hence we only need to compute the function J(z) in eq. (78) at the point of the mean field solution $z = Z_{\rm MF}(\lambda) = \frac{1}{2} \left(1 - \sqrt{1 - 4\lambda}\right)$.

The integral J(z) can be rewritten using the elastic generating function C(t) defined in (91) for an arbitrary elastic kernel ϵ_k as

$$J(z) = \int_0^\infty dt \, \mathcal{C}(t) \, \left[e^{2tz} \left(tz^2 + z \right) - z - 3tz^2 \right] \, . \tag{102}$$

4.4.2 Standard elasticity

Standard elasticity has the form $e(p) = p^2 + 1$ and using (94) one finds:

$$J(z) = \frac{1}{2}z \left[2z + (1 - 3z)\log(1 - 2z)\right] .$$
(103)

This result is in agreement with eq. (150) of [10]. The value of α to be used in (101) for extrapolation to d = 1 (i.e. $\epsilon = 3$) is $\alpha = -2/3$.

4.4.3 Flat contact line and generalization

From the form $e(p) = \sqrt{p^2 + (1-a)^2} + a$ and using (99) one finds

$$J(z) = z \left[\frac{2z(a+z-3az)}{1-2z} + (a-3z)\log(1-2z) \right].$$
(104)

This agrees with (E10) of [10] for a = 0 and with (103) up to a global factor of 2, as predicted in [10]. The value of α to be used in (101) for extrapolation to d = 1 (i.e. $\epsilon = 1$) is $\alpha = -2/9$.

4.4.4 Contact line for arbitrary angles θ and φ

Having tested our general formula on the known cases we can now apply them to the case of the elastic kernel (62). Let us specify the relevant parameters and functions. For the contact line $\beta = 1$, $d_c = 2$, and

$$m^{2} = \mu = \epsilon_{q=0} = \frac{\gamma \kappa}{t} \sin \theta \cos \varphi \qquad (105)$$

$$K = \gamma \sin^2 \theta \tag{106}$$

$$\kappa = \frac{1}{L_c} = \sqrt{\frac{\rho g}{\gamma}} \quad , \quad t = \sqrt{\frac{\sin(\theta + \varphi) + 1}{2}} \tag{107}$$

$$\tilde{\epsilon}_q = e(p) = \tilde{e}(r) = 1 + \frac{(r^2 - 1) [t(r+t) + 1] \sin \theta}{(r^2 + 3rt + 3t^2 - 1) \cos \varphi}$$
(108)

$$r = \sqrt{1 + \frac{q^2}{\kappa^2}} = \sqrt{1 + \frac{\cos^2 \varphi}{t^2 \sin^2 \theta} p^2} , \quad p = \frac{K}{\mu} q .$$
 (109)

We have defined for convenience a new function $\tilde{e}(r)$ since r is the implicit variable in the final formula. Note that for the flat interface $\theta + \varphi = \pi/2$, one recovers $\tilde{e}(r) = r$.

The generating function $\tilde{Z}(\lambda)$ for the contact line can thus be computed from (101) using (85) and performing the integral in dimension $d_c = 2$. Upon a change of variable we get:

$$J(z) = \frac{t^2 \sin^2 \theta}{\cos^2 \varphi}$$

$$\times \int_{1}^{\infty} \mathrm{d}r \, r \left[\frac{z^2}{[\tilde{e}(r) - 2z]^2} + \frac{z}{\tilde{e}(r) - 2z} - \frac{z}{\tilde{e}(r)} - 3\frac{z^2}{\tilde{e}(r)^2} \right].$$
(110)

Performing this integral analytically, or even the one involved in computing C(t), is rather awkward, and we have preferred to use numerical integration.

To relate to the experiment of [1], we have computed J(z) numerically for $\theta = 40^{\circ}$, $\varphi = 0^{\circ}$. We give some explicit values: J(-1/2) = -0.493277, J(-1) = -3.10539, J(-3/2) = -8.79405, J(-2) = -18.1329. From this we have computed and plotted $\tilde{Z}(\lambda)$ in figure 9, using the extrapolation to $\epsilon = 1$ in the one-loop result, i.e. using (101) with $\alpha = -2/9$. This is compared to the result of the scaled kernel $e(p) = \sqrt{p^2 + 1}$. The experimental data are also plotted on figure 9. Note that using the correct elasticity allows to get closer to the experimental data than using $e(p) = \sqrt{p^2 + 1}$, which is the kernel for the flat interface, and the only one previously available in the literature.

4.5 Local statistics of avalanches

4.5.1 Basic definitions

It is also possible to study in experiments the statistics of the avalanches occurring within a given portion of the elastic object, e.g. of the contact line. One thus defines the size of a local avalanche, weighted by a characteristic



Fig. 9. The generating function $\tilde{Z}(\lambda)$ of avalanche sizes as defined in (75). The curves are from bottom to top: Red/solid: mean-field result. Black dots: The experimental data of [1]. Dashed/green: Eq. (110) integrated numerically with the correct elasticity kernel (109) for $\theta = 40^{\circ}$, $\varphi = 0^{\circ}$. Dot-Dashed/orange: The same with the elastic kernel for $\theta = 90^{\circ}$, $\varphi = 0^{\circ}$, $\tilde{\epsilon}_q = \sqrt{1+q^2/\kappa^2}$.

function $\phi(x)$, e.g. a δ -distribution localized at x = 0, as ⁴:

$$S^{\phi} := \int_{0}^{L} \mathrm{d}^{d} x \, \phi(x) \left[u_{w}^{+}(x) - u_{w}^{-}(x) \right]$$
$$= \int_{0}^{L} \mathrm{d}^{d} x \, \phi(x) \left[h_{w}^{+}(x) - h_{w}^{-}(x) \right] \,. \tag{111}$$

In [10], we have shown how the characteristic function

$$Z^{\phi}(\lambda) := \frac{1}{\langle S^{\phi} \rangle} \left\langle e^{\lambda S^{\phi}} - 1 \right\rangle$$
(112)

as well as the distribution of sizes can be computed within mean-field theory, valid at and above the upper critical dimension d_c . Since it already required some quite involved instanton calculus, we did not attempt to perform the ϵ expansion. Here we will perform the ϵ expansion, to one loop $\mathcal{O}(\epsilon)$, not on the full distribution, but on a simpler quantity, the second moment. More precisely, we will compute the universal amplitude ratio between the second global and local moments, quantities easily accessible in experiments and numerics. In fact, it was already measured in [13] in the case of standard local elasticity and the present analytical result was quoted. Here we provide the details of the calculation and predict this ratio for the contact line which has not yet been measured.

4.5.2 Universal amplitude ratios between global and local avalanches: general formulation

Analogous to (71), we can define the scale of local avalanches as

$$S_m^{\phi} \coloneqq \frac{\langle (S^{\phi})^2 \rangle}{2 \langle S^{\phi} \rangle} . \tag{113}$$

From Eq. (F12) and (196) in [10] one has the general exact relation:

$$S_m^{\phi} = -\frac{1}{\int_x \phi(x)} \partial_w \int_{xyzz'} \phi(y)\phi(x)g_{xz}g_{yz'}\Delta_{zz'}[w]\Big|_{w=0^+} ,$$

where $g_{xy} = \int_q e^{iq(x-y)} \epsilon_q^{-1}$ and $\Delta_{zz'}[w] = -R''_{zz'}[w]$ is the functional derivative taken at a uniform $w_t = w$ of the functional R[w] (see e.g. [14,10] for its definition), the limit $w = 0^+$ being taken at the end. For $\phi(x) = 1$, using $\int_{zz'} \Delta_{zz'}[w] = L^d \Delta'(w)$ one recovers

$$S_m = \frac{\langle S^2 \rangle}{2 \langle S \rangle} = -m^{-4} \Delta'(0^+) . \qquad (114)$$

Here we use the notations of section (4.2.2) so that $m^2 = \epsilon_{k=0}$.

We can now use the results of the ϵ -expansion [15, 16, 17], or its first-principle derivation in formula (463) of [18]:

$$\Delta_{zz'}[w] = \delta_{zz'}\Delta(w) - (g_{zz'}^2 - \delta_{zz'}\int_t g_t^2)\Delta'(w)^2 + \mathcal{O}(\epsilon^3) .$$

Hence one obtains the ratio

$$a_{\phi} := \frac{S_{m}^{\phi}}{S_{m}}$$
(115)
$$= \frac{m^{4}}{\int_{x} \phi(x)} \int_{xyz} \phi(y)\phi(x)g_{xz}g_{yz} -2m^{4}\Delta''(0^{+})\frac{1}{\int_{x} \phi(x)} \times \int_{xyzz'} \phi(y)\phi(x)g_{xz}g_{xz'}(g_{zz'}^{2} - \delta_{zz'}\int_{t} g_{t}^{2}) + \mathcal{O}(\epsilon^{2}) ,$$

a formula valid for any function $\phi(x)$.

We know study avalanches which occur on a given subspace of co-dimension d', defined by $x_{\perp} = 0$, i.e. of dimension $d_{\phi} = d - d'$, of the *d*-dimensional manifold, i.e. we choose

$$\phi(x) = \eta \delta^{d'}(x_{\perp}) . \tag{116}$$

 η will be specified later. The above formula leads to

$$a_{\phi} = m^{4} \eta I_{2}(d') + 2m^{4} \eta \left[I_{2}(d) I_{2}(d') - I_{A}(d,d') \right] \Delta''(0) + \dots$$

$$= m^{4} \eta I_{2}(d') \left[1 + \frac{2\tilde{\Delta}''(0)}{\epsilon} \left(1 - \frac{I_{A}(d,d')}{I_{2}(d)I_{2}(d')} \right) \right] + \dots$$
(117)

using the definition (83) of the rescaled disorder, $\tilde{\Delta}''(0) = \Delta''(0)\epsilon I_2(d)$, and defining the integrals

$$I_2(d) := \checkmark = \int \frac{\mathrm{d}^d q}{(2\pi)^d} \frac{1}{\epsilon_q^2} \qquad (118)$$

$$I_A(d, d') := \bigwedge_{=}^{dd' k} \int \frac{d^{d'}k}{(2\pi)^{d'}} \frac{d^d q}{(2\pi)^d} \frac{1}{\epsilon_k^2} \frac{1}{\epsilon_q} \frac{1}{\epsilon_{k+q}} .$$
(119)

⁴ Note that the variable $\phi(x)$ entering in S^{ϕ} and a^{ϕ} bears no relation to the plate angle φ introduced previously.

Note that in the last integral the momentum in the upper bubble runs over a space of dimension d' < d.

To perform the ϵ expansion we now use the scaling form (81) of the elastic kernel. This yields to zeroth and first order in $\epsilon = d_c - d$:

$$a_{\phi} = \eta \mu^{d'} K^{-d'/\beta} \tilde{I}_{2}(d') \left[1 + \frac{2\tilde{\Delta}''(0)}{\epsilon} \left(1 - \frac{\tilde{I}_{A}(d,d')}{\tilde{I}_{2}(d)\tilde{I}_{2}(d')} \right) \right] + \cdots$$
(120)

in terms of the dimensionless integrals

$$\tilde{I}_2(d) := \checkmark = \int \frac{\mathrm{d}^d q}{(2\pi)^d} \frac{1}{e(q)^2} \tag{121}$$

$$\tilde{I}_A(d,d') := - \int \frac{\mathrm{d}^{d'}k}{(2\pi)^{d'}} \frac{\mathrm{d}^d q}{(2\pi)^d} \frac{1}{e(k)^2} \frac{1}{e(q)} \frac{1}{e(k+q)} . \quad (122)$$

To obtain a meaningful amplitude we now make the choice

$$\eta = \left(\frac{K^{1/\beta}}{\mu}\right)^{d'} \quad . \tag{123}$$

Note that comparing with experiments then requires a precise knowledge of the internal length scale $K^{1/\beta}/\mu$ defined by the (renormalized) mass $m^2 = \mu^\beta$ and elastic constant $K = \lim_{q \gg \kappa} \epsilon_q/q^\beta$. Alternatively, it may be used as a method to measure this length.

4.5.3 Standard local elasticity

We know specify to standard elasticity $\epsilon_k = Kk^2 + m^2$, i.e. $\beta = 2$, $d_c = 4$ and $e(p) = p^2 + 1$. We study d' = 1.

We need the integrals:

$$\tilde{I}_2(d) = \Gamma\left(2 - \frac{d}{2}\right) (4\pi)^{-d/2}$$
 (124)

$$\tilde{I}_2(d'=1) = \int \frac{dq}{2\pi} \frac{1}{(q^2+1)^2} = \frac{1}{4}$$
(125)

$$\frac{I_A(d,d')}{\tilde{I}_2(d)\tilde{I}_2(d')} - 1 = \frac{I}{\tilde{I}_2(d)\tilde{I}_2(d')}$$
(126)

This involves the integral

$$I = \int \frac{dk}{2\pi} \frac{1}{(k^2 + 1)^2} \int \frac{dq}{2\pi} \frac{d^{d-1}p}{(2\pi)^d}$$
(127)
 $\times \left[\frac{1}{[(k+q)^2 + p^2 + 1](q^2 + p^2 + 1)} - \frac{1}{(q^2 + p^2 + 1)^2} \right]$

The calculation of this integral is performed in Appendix A. To the order ϵ we are working, we only need to compute it in $d = d_c = 4$. We find

$$I = (4\pi)^{-\frac{5}{2}} \Gamma\left(\frac{3}{2}\right) \left[4 + \frac{2}{9}(7\sqrt{3} - 18)\pi\right] .$$
 (128)

Putting all these results together we obtain

$$a_{\phi} = \frac{1}{4} + \alpha \left(1 - \pi + \frac{7\pi}{6\sqrt{3}} \right) + \mathcal{O}(\epsilon^2)$$

= 0.25 - 0.0254934\alpha + \mathcal{O}(\epsilon^2) (129)

with $\alpha = -\frac{\epsilon}{3}(1-\zeta_1)$. This is the expansion for a manifold with co-dimension one in $\epsilon = 4 - d$.

For the 1-dimensional string, simulated in [13], we need to extrapolate to $\epsilon = 3$ which yields $\alpha = -2/3$. The two possible Padé expansions are

$$\frac{1}{a_{\phi}} = \frac{1}{\frac{1}{4} + 0.0169956} = 3.74538 , \qquad (130)$$

$$\frac{1}{a_{\phi}} = 4 - 16\alpha \left[1 + \left(\frac{7}{6\sqrt{3}} - 1\right)\pi \right] = 3.72807 .$$
(131)

Hence an estimate of the ratio is

$$\frac{1}{a_{\phi}} = 3.74 \pm 0.01 \ . \tag{132}$$

Another expansion is the expansion in fixed dimension in powers of $\tilde{\Delta}$. It is likely to be less precise than the ϵ -expansion, but being simpler in spirit we indicate here its prediction. One evaluates directly the integrals in d =d' = 1. Using the results from appendix A, this yields

$$a_{\phi} = \frac{1}{4} \left[1 + \frac{2}{9} \frac{\tilde{\Delta}''(0^+)}{\epsilon} + \dots \right] ,$$
 (133)

and inserting the one-loop value $\tilde{\Delta}''(0^+) = -\alpha = \frac{2}{9}\epsilon$ we find the estimate $1/a_{\phi} \approx 3.81$, slightly larger than (132).

4.5.4 Contact-line elasticity

For the contact-line elasticity, the $\epsilon = 2 - d$ expansion can be done as indicated above. As it requires performing two-dimensional integrals we will only give here the fixeddimension estimate for d = d' = 1, which requires only a one-dimensional integral on each momentum. We further perform them numerically using the form (109). We give two examples:

(i) flat interface $\varphi + \theta = \pi/2$. Then $e(p) = \sqrt{p^2 + 1}$ and we obtain

$$\tilde{I}_2(1) = \frac{1}{2} \tag{134}$$

$$2\left[1 - \frac{\tilde{I}_A(1,1)}{\tilde{I}_2(1)\tilde{I}_2(1)}\right] \approx 0.312811\dots$$
(135)

This yields

$$a_{\phi} = \frac{1}{2} \left(1 + 0.312811 \frac{2}{9} + \dots \right)$$
(136)

and $1/a_{\phi} \approx 1.87$.

(ii) experiments of Ref. [1], i.e. $\varphi = 0^{\circ}, \theta = 40^{\circ}$: We find

$$I_{2}(1) \approx 0.41867$$
(137)
$$2 \left[1 - \frac{\tilde{I}_{A}(1,1)}{\tilde{I}_{2}(1)\tilde{I}_{2}(1)} \right] \approx 0.333101 \cdots$$
(138)

This yields

$$a_{\phi} = 0.41867 \left(1 + 0.333101 \frac{2}{9} + \ldots \right)$$
 (139)

and $1/a_{\phi} \approx 2.22$.

It is interesting to note that the "elastic length" $\eta =$ K/μ which enters the definition (111), (116) of S^{ϕ} and a_{ϕ} reads:

$$\eta = \frac{K}{\mu} = L_c \frac{\sin\theta}{\cos\varphi} \sqrt{\frac{\sin(\theta + \varphi) + 1}{2}}$$
(140)

i.e., it only involves the capillary length and geometric prefactors, thus is well-known experimentally. In addition, one can measure the dependence of a_{ϕ} on the plate angle φ .

Acknowledgments

It is a pleasure to thank Etienne Rolley and Alberto Rosso for stimulating discussions.



for standard elasticity.

References

- 1. P. Le Doussal, K.J. Wiese, S. Moulinet and E. Rolley, Height fluctuations of a contact line: a direct measurement of the renormalized disorder correlator, arXiv:0904.1123 (2009).
- 2. J.F. Joanny and P.G. De Gennes, A model for contactangle hysteresis, J. Chem. Phys. 81 (1984) 552-562.
- 3. Y. Pomeau and J. Vannimenus, Contact-angle on heterongenous surfaces - weak heterogeneities, Journal of Colloid and Interface Science 104 (1985) 477-488.
- 4. V. S. Nikolayev and D. A. Beysens, Equation of motion of the triple contact line along an inhomogeneous surface, Europhys. Lett. 64 (2003) 763.
- 5. F. Brochard and P. G. De Gennes, Collective modes of a $contact\ line,\ Langmuir\ 7\ (1991)\ 3216–3218.$
- 6. A. Checco, Non-linear elasticity of a liquid contact line, EPL 85 (2009) 16002.
- 7. R. Golestanian and E. Raphael, Relaxation of a moving contact line and the landau-levich effect, Europhys. Lett. **55** (2001) 228–34.
- 8. P. Le Doussal, K.J. Wiese, E. Raphael and Ramin Golestanian, Can non-linear elasticity explain contact-line roughness at depinning?, Phys. Rev. Lett. 96 (2006) 015702, cond-mat/0411652.
- 9. C. Bachas, P. Le Doussal and K.J. Wiese, Wetting and minimal surfaces, Phys. Rev. E 75 (2007) 031601, hepth/0606247.
- 10. P. Le Doussal and K.J. Wiese, Size distributions of shocks and static avalanches from the functional renormalization group, arXiv:0812.1893 (2008).
- 11. K.J. Wiese and P. Le Doussal, Functional renormalization for disordered systems: Basic recipes and gourmet dishes, Markov Processes Relat. Fields 13 (2007) 777–818, condmat/0611346.
- 12. P. Le Doussal, A.A. Middleton and K.J. Wiese, Statistics of static avalanches in a random pinning landscape, arXiv:0803.1142 (2008).
- 13. A. Rosso, P. Le Doussal and K.J. Wiese, Avalanche-size distribution at the depinning transition: A numerical test of the theory, arXiv:0904.1123 (2009).
- 14. P. Le Doussal, Finite temperature Functional RG, droplets and decaying Burgers turbulence, Europhys. Lett. 76 (2006) 457-463, cond-mat/0605490.
- 15. P. Chauve, P. Le Doussal and K.J. Wiese, Renormalization of pinned elastic systems: How does it work beyond one loop?, Phys. Rev. Lett. 86 (2001) 1785-1788, condmat/0006056.
- 16. P. Le Doussal, K.J. Wiese and P. Chauve, 2-loop functional renormalization group analysis of the depinning transition, Phys. Rev. B 66 (2002) 174201, cond-mat/0205108.
- 17. P. Le Doussal, K.J. Wiese and P. Chauve, Functional renormalization group and the field theory of disordered elastic systems, Phys. Rev. E (2004) 026112, cont-mat/ 0304614.
- 18. P. Le Doussal, Exact results and open questions in first principle functional RG, arXiv:0809.1192 (2008).

Pierre Le Doussal, Kay Jörg Wiese: Elasticity of a contact-line and avalanche-size distribution at depinning 15

A Standard elasticity: Calculation of an integral

The integral defined in the text can be written as

$$I = \int \frac{dk}{2\pi} \int \frac{dq}{2\pi} \frac{d^{d-1}p}{(2\pi)^d} \int_{t,t_1,t_2>0} te^{-t(k^2+1)} \left[e^{-t_1((k+q)^2+p^2+1)+t_2(q^2+p^2+1)} - e^{-(t_1+t_2)(q^2+p^2+1)} \right]$$

= $(4\pi)^{-\frac{d+1}{2}} \int_{t,t_1,t_2>0} t(t_1+t_2)^{-\frac{d-1}{2}} \left[(tt_1+tt_2+t_1t_2)^{-1/2} - (t(t_1+t_2))^{-1/2} \right] e^{-t-t_1-t_2} ,$ (141)

using the general formula $\int_q e^{-q \cdot C \cdot q} = (4\pi)^{-d/2} (\det C)^{-1/2}$. Performing the changes of variables $t_1 \to tt_1$ and $t_2 \to tt_2$ yields

$$I = (4\pi)^{-\frac{d+1}{2}} \int_0^\infty dt \int_0^\infty dt_1 \int_0^\infty dt_2 \, e^{-t(t_1+t_2+1)} t^{\frac{5}{2}-\frac{d}{2}} \left(t_1+t_2\right)^{\frac{1}{2}-\frac{d}{2}} \left[\frac{1}{\sqrt{t_2t_1+t_1+t_2}} - \frac{1}{\sqrt{t_1+t_2}}\right] \,. \tag{142}$$

Integration over t yields

$$I = (4\pi)^{-\frac{d+1}{2}} \int_0^\infty \mathrm{d}t_1 \int_0^\infty \mathrm{d}t_2 \left(t_1 + t_2\right)^{\frac{1}{2} - \frac{d}{2}} \left(t_1 + t_2 + 1\right)^{\frac{d-7}{2}} \Gamma\left(\frac{7}{2} - \frac{d}{2}\right) \left[\frac{1}{\sqrt{t_2 t_1 + t_1 + t_2}} - \frac{1}{\sqrt{t_1 + t_2}}\right] .$$
(143)

Setting $t_1 = st$, $t_2 = (1 - s)t$, one obtains

$$I = (4\pi)^{-\frac{d+1}{2}} \int_0^\infty dt \int_0^1 ds \, t^{1-\frac{d}{2}} (t+1)^{\frac{d-7}{2}} \Gamma\left(\frac{7}{2} - \frac{d}{2}\right) \left[\frac{1}{\sqrt{1+(1-s)st}} - 1\right]$$
$$= (4\pi)^{-\frac{d+1}{2}} \int_0^\infty dt \, t^{1-\frac{d}{2}} (t+1)^{\frac{d-7}{2}} \Gamma\left(\frac{7}{2} - \frac{d}{2}\right) \left[\frac{2}{\sqrt{t}} \operatorname{arccot}\left(\frac{2}{\sqrt{t}}\right) - 1\right] .$$
(144)

One can perform the integral directly in d = 1, with the result

$$I = -\frac{1}{144} \ . \tag{145}$$

Thanks to the counterterm it also admits a limit for d = 4:

$$I = \frac{1}{8\pi^2} \left[\frac{1}{2} + \frac{\pi}{36} (-18 + 7\sqrt{3}) \right] .$$
 (146)

Finally the expression for any d is

$$I = \frac{9\sqrt{3}2^{d}\Gamma\left(2-\frac{d}{2}\right){}_{3}F_{2}\left(\frac{1}{2},1,2-\frac{d}{2};-\frac{1}{2},\frac{3}{2};\frac{1}{4}\right) + 2\sqrt{\pi}\left(9\sqrt{3}2^{d}-43^{d/2}(d+3)\right)\Gamma\left(\frac{3}{2}-\frac{d}{2}\right)}{4^{d+1}\pi^{d/2}9\sqrt{3}} - 2^{-2-d}\pi^{-d/2}\Gamma\left(2-\frac{d}{2}\right) .$$
(147)

А

The combination $2\left[1 - \frac{\tilde{I}_A(d,1)}{\tilde{I}_2(1)\tilde{I}_2(d)}\right]$ is plotted on Fig. 10.

Contents

1	Intre	Introduction $\ldots \ldots 1$			
2	Mod	lel	2		
3	Con	tact-line elasticity	3		
	3.1	Model in shifted coordinates	3		
	3.2	Zero mode and calculation of the mass	3		
	3.3	Elastic energy for arbitrary deformations	4		
	3.4	The contact-line profile as a means of mea-			
		suring ϵ_q	$\overline{7}$		
4	Ava	lanche-size distributions	$\overline{7}$		
	4.1	The model	$\overline{7}$		
	4.2	Global statistics of avalanches	8		
		$4.2.1 \text{Definitions} \dots \dots \dots \dots \dots \dots \dots \dots \dots $	8		
		4.2.2 Results from one-loop FRG	9		
	4.3	From $\tilde{Z}(\lambda)$ to $p(s)$	10		
		$4.3.1 \text{General formalism} \dots \dots \dots \dots$	10		
		$4.3.2 \text{Special cases} \dots \dots \dots \dots \dots \dots \dots \dots \dots $	10		

	Standa	ard local elasticity:	10
	Elastic	city of flat contact line and general-	
		ization:	10
4.4	More of	on the generating function $\tilde{Z}(\lambda)$	11
	4.4.1	Generating function in the general	
		case	11
	4.4.2	Standard elasticity	11
	4.4.3	Flat contact line and generalization	11
	4.4.4	Contact line for arbitrary angles θ	
		and φ	11
4.5	Local	statistics of avalanches	11
	4.5.1	Basic definitions	11
	4.5.2	Universal amplitude ratios between	
		global and local avalanches: general	
		formulation \ldots \ldots \ldots \ldots \ldots \ldots	12
	4.5.3	Standard local elasticity	13
	4.5.4	Contact-line elasticity	13
Stan	idard el	asticity: Calculation of an integral	15