

Statics and dynamics of elastic manifolds in media with long-range correlated disorder

Andrei A. Fedorenko, Pierre Le Doussal, and Kay Jörg Wiese

CNRS-Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, 24 rue Lhomond, 75231 Paris, France

(Dated: September 10, 2006)

We study the statics and dynamics of an elastic manifold in a disordered medium with quenched defects correlated as $\sim r^{-a}$ for large separation r . We derive the functional renormalization group equations to one-loop order which allow to describe the universal properties of the system in equilibrium and at the depinning transition. Using a double $\varepsilon = 4 - d$ and $\delta = 4 - a$ expansion we compute the fixed points characterizing different universality classes and analyze their regions of stability. The long-range disorder-correlator remains analytic but generates short-range disorder whose correlator exhibits the usual cusp. The critical exponents and universal amplitudes are computed to first order in ε and δ at the fixed points. At depinning a velocity-versus-force exponent β larger than unity can occur. We discuss possible realizations using extended defects.

I. INTRODUCTION

Elastic objects in random media are the simplest example of disordered system exhibiting metastability, glassy behavior and dimensional reduction, which are difficulties present in a broader class of disordered systems [1, 2, 3]. They can be used to model a remarkable set of experimental systems. Domain walls in magnets behave as elastic interfaces and can experience either random bond disorder (RB) as in ferromagnets with non-magnetic impurities or random-field disorder (RF) as in disordered antiferromagnets in an external magnetic field [4]. The interface between two immiscible liquids in a porous medium exhibits the same behavior and undergoes a depinning transition as the pressure difference is increased [5]. Charge density waves (CDW) in solids show a similar conduction threshold [6]. Another example of periodic systems are vortex lines in superconductors which can form different glass phases in the presence of weak disorder [7, 8, 9]. In all these systems the interplay between elastic forces which tend to keep the system ordered, i.e. flat or periodic, and quenched disorder, which promotes deformations of the local structure, forms a complicated energy landscape with numerous metastable states. This results in glassy properties and a nontrivial response of the system to external perturbations. In particular, the interface becomes rough with displacements growing with the distance x as

$$C(x) \sim x^{2\zeta}, \quad (1)$$

where ζ is the roughness exponent. Elastic periodic structures in the presence of disorder lose their strict translational order and form quasi long-range order characterized by a slow growth of displacements,

$$C(x) = \mathcal{A}_d \ln |x| \quad (2)$$

where the amplitude \mathcal{A}_d is universal in the simplest case. At zero temperature, a driving force f exceeding the threshold value f_c is required to set the elastic manifold into steady motion with a velocity v which vanishes as $v \sim (f - f_c)^\beta$ at the transition point. The correlation length diverges close to the transition $f = f_c$ as $\xi \sim (f - f_c)^{-\nu}$ and the characteristic time as $\tau \sim \xi^z$, where z is the dynamic critical exponent. Note that the roughness exponent and the universal amplitudes

determined at the depinning transition are in general different from the exponent and amplitudes measured in equilibrium.

Two methods were developed to study the statics of an elastic manifold in a disordered medium. One of them is the Gaussian variational approximation performed in replica space, which can be applied to both classes of elastic manifolds, i.e. to interfaces [10] and to periodic systems [9, 11]. Within this approach, which is believed to be exact in the mean-field limit, i.e. when the manifold lives in a space of infinite dimensions, metastability is described by breaking of replica symmetry, that allows one to compute the static correlation functions and to obtain different thermodynamic properties. Another method which can be applied to dynamics as well as to statics is the functional renormalization group (FRG) [12]. Simple scaling arguments show that large-scale properties of elastic systems are governed by disorder for $d < d_{uc} = 4$ and that perturbation theory in the disorder breaks down on scales larger than the so-called Larkin scale [13]. To overcome this difficulty one performs a renormalization-group analysis. It was shown that in this case one has to renormalize the whole disorder correlator which becomes a non-analytic function beyond the Larkin scale [12, 14, 15, 16]. The appearance of a non-analyticity in the form of a cusp at the origin is related to metastability, and nicely accounts for the generation of a threshold force at the depinning transition. It was recently shown that the FRG can unambiguously be extended to higher loop order so that the underlining non-analytic field theory is probably renormalizable to all orders [17, 18, 19]. Although the two methods are very different, they provide a fairly consistent picture of the statics, and recently a relation between them was found [20]. There is also good agreement with results of numerical simulations, not only for critical exponents [21, 22, 23], but also for the whole renormalized disorder correlator [24]. However, many questions remain open. Although the dynamics in the vicinity of the depinning transition and at zero temperature is well understood, there is no satisfactory theory for finite temperature, and in particular for the thermal rounding of the depinning transition [25]. It is also remarkable that the exponent β in experiments on depinning is usually larger than 1, while FRG and numerical simulations of elastic systems with weak disorder give values smaller than 1.

Most studies of elastic manifolds in a disordered medium

treat uncorrelated point-like disorder. Real systems, however, often contain extended defects in the form of linear dislocations, planar grain boundaries, three-dimensional cavities, etc. It is known that such extended defects, or point-like defects with sufficiently long-range correlations can change the bulk critical behavior [26, 27, 28, 29, 30, 31, 32]. Flux lines in superconductors are the most prominent example. The pinning of the flux lines by disorder prevents the dissipation of energy and determines the critical current J_c , which is of great importance for applications. It was found that extended defects produced, for instance, by heavy ion irradiation, can increase J_c by several orders of magnitude [33]. Systems with anisotropic orientation of extended defects can be described by a model in which all defects are strongly correlated in ε_d dimensions and randomly distributed over the remaining $d - \varepsilon_d$ dimensions. The case $\varepsilon_d = 0$ is associated with uncorrelated point-like defects, while extended columnar or planar defects are related to the cases $\varepsilon_d = 1$ and 2 respectively. The bulk-critical behavior in the presence of this type of disorder was studied in Refs. [26, 27, 28, 29] using a perturbative RG analysis in conjunction with a double expansion in $\varepsilon = 4 - d$ and ε_d . The pinning of flux lines by columnar disorder was studied in Ref. [34], where it was shown that the system forms a Bose glass phase with flux lines strongly localized on the columnar defects, resulting in a zero dc linear resistivity. It was argued recently that the topologically ordered glass phase (Bragg glass) formed by flux lines can be destroyed in the vicinity of a single planar defect [35]. It has been shown that the small dispersion in orientation of columnar defects forms a new thermodynamic phase called “splayed glas” [36]. In this phase, the entanglement of flux lines enhances significantly the transport of superconductors [37]. Competition between various types of disorder, point and columnar, has also been studied, at equilibrium [38, 39] and in the moving phases [40].

In the case of an isotropic distribution of disorder, power-law correlations are the simplest example with the possibility for a scaling behavior with new fixed points (FPs) and new critical exponents. The bulk-critical behavior of systems in which defects are correlated according to a power-law r^{-a} for large separation r was studied in Refs. [30, 31, 32]. The power-law correlation of defects in d -dimensional space with exponent $a = d - \varepsilon_d$ can be ascribed to randomly distributed extended defects of internal dimension ε_d with random orientation. For example, $a = d$ corresponds to uncorrelated point-like defects, $a = d - 1$ ($a = d - 2$) describes infinite lines (planes) of defects with random orientation. In general one would probably not expect a pure power-law decay of correlations. However, if the correlations of defects arise from different sources with a broad distribution of characteristic lengthscales, one can expect that the resulting correlations will over several decades be approximated by an effective power-law [30]. If the correlation function of disorder can be expressed as a finite sum of power-law contributions $\sum_i c_i r^{-a_i}$, one can expect that the scaling behavior is dominated by the term with the smallest a_i [30]. Power-law correlations with a non-integer value $a = d - d_f$ can be found in systems containing defects with fractal dimension d_f [41]. For example, the behavior of ^4He in aerogels is argued to be

described by an XY model with LR correlated defects [42]. This is closely related to the behavior of nematic liquid crystals enclosed in a single pore of aerosil gel which was recently studied in Ref. [43], using the approximation in which the pore hull is considered a disconnected fractal. Finally, studies of the Kardar-Parisi-Zhang (KPZ) equation with power-law correlations in time [44] bear connections to the case $d = 1$ considered here. However the perturbative method used there cannot address directly the zero temperature (strong KPZ coupling) phase, contrarily to our present study.

In the present paper we study the statics and dynamics of elastic manifolds in the presence of (power law) LR correlated disorder using the FRG approach to one-loop order. The paper is organized as follows. Section II introduces the model. Possible physical realizations are considered in Section III. Section IV describes the dynamical formalism and perturbation theory. In Section V we renormalize the theory and derive the FRG equations to one-loop order. In Section VI we study random bond, in Section VII random field and in Section VIII periodic disorder. In Section IX we discuss fully isotropic extended defects. In the final Section we summarize the obtained results and our conclusions.

II. THE MODEL

We consider a d -dimensional elastic manifold embedded in a D -dimensional space with quenched disorder. The configuration of the manifold is described by a N -component displacement field denoted below $u(x)$, or equivalently u_x , where x denotes the d -dimensional internal coordinate of the manifold. For example, a domain wall corresponds to $d = D - 1$ and $N = 1$, a CDW to $d = D$ and $N = 1$, and a flux lattice to $d = D$ and $N = 2$. In what follows, we focus for simplicity on the case $N = 1$ and elastic objects with short-range elasticity. Extensions to $N > 1$ and LR elasticity are straightforward for the statics. The energy of the manifold in presence of disorder is defined by the Hamiltonian

$$\mathcal{H} = \int d^d x \left[\frac{c}{2} (\nabla u(x))^2 + V(x, u(x)) \right], \quad (3)$$

where c is the elasticity and V a random potential. In this paper we study the model where the second cumulant of the random potential has the form:

$$\overline{V(x, u)V(x', u')} = R_1(u - u')\delta^d(x - x') + R_2(u - u')g(x - x'). \quad (4)$$

The first part corresponds to point-like disorder with short-range (SR) correlations in internal space. The second part corresponds to long-range (LR) disorder in internal space and the function $g(x) \sim x^{-a}$ at large x . For convenience we normalize it so that its Fourier transform is $g(q) = |q|^{a-d}$ at small q with unit amplitude. A priori we are interested in the case $a < d$ where the correlations decay sufficiently slowly in internal space. We denote everywhere below $\int_q = \int \frac{d^d q}{(2\pi)^d}$ and $\int_x = \int d^d x$. The short-scale UV cutoff is implied at $q \sim \Lambda$ and the size of the system is L .

One could start with model (4), setting $R_1 = 0$; however as we show below a non-zero R_1 is generated under coarse graining. Note that the functions $R_i(u)$ can themselves a priori be SR, LR, or periodic in the direction of the displacement field u . For SR disorder in internal space only, i.e. $R_2 = 0$, these cases are usually referred to as random bond (RB), random field ($R_1(u) \sim |u|$ at large u) (RF) and random periodic (RP) universality classes. Below we discuss how these classes extend to the case of LR internal disorder (R_2 non zero).

The model (3) and (4) could easily be studied using presently available numerical algorithms for directed manifolds, in its statics (e.g. exact ground state determinations) and its dynamics (e.g. critical configuration at depinning), by directly implementing a random potential correlated as described by (4). It is also interesting to examine which type of correlations in a random medium can naturally lead to (4) and how such disorder could be realized from e.g. distributions of extended defects, since some of them may be experimentally feasible.

III. REALIZATIONS AND UNIVERSALITY CLASSES

A. Defect potential

Let us first recall how long-range correlations can arise in the potential created by defects. To this purpose call $v(r)$ the defect potential, in the simplest case taken to be proportional to defect density. Consider for simplicity a large number of weak defect lines with a uniform and isotropic distribution in a space of dimension D . These create an almost Gaussian random potential $v(r)$ with:

$$\overline{v(r)v(r')} \sim \frac{v_{\text{LR}}^2}{|r - r'|^a} \quad \text{for } r \rightarrow \infty \quad (5)$$

and $a = D - 1$. To derive this, consider defects of finite radius a_d . The probability that point r' is contained in the defect going through r is $\sim (a_d/|r - r'|)^{D-1}$, i.e. inversely proportional to the sphere of radius $|r - r'|$. This is easily generalized to isotropic distributions of extended defects of internal dimension ε_d , with $a = D - \varepsilon_d$. Note that by extended defects we mean defects which are perfectly correlated along their internal dimension. Generalizations where defects are themselves (anisotropic) fractals can also be considered.

An important case is a uniform distribution of extended defects in D dimensional space, but isotropic only within a linear subspace of dimension D' . For instance one can irradiate a material in the bulk while simultaneously rotating it along an axis. This produces a distribution of linear defects ($\varepsilon_d = 1$), isotropic within the plane ($D' = 2$), and normal to the axis (see Fig. 1). More generally this yields a defect potential with second cumulant

$$\overline{v(r,z)v(r',z)} = g(r - r')f(z - z') \quad (6)$$

$$g(r) \sim r^{-a},$$

while $f(z)$ is short-ranged (here $r \in R^{D'}$, $z \in R^{D-D'}$, $a = D' - \varepsilon_d$).




FIG. 1: Linear defects randomly and isotropically distributed on parallel planes with random distances between them. This geometry mimics distribution (6).

Although we mostly discuss extended defects, other sources of long-range correlations are possible, such as defects where each single one creates a long-ranged disorder potential, or a substrate matrix itself quenched at a critical point.

B. Coupling to the manifold

We now examine how the long-range correlated defect potential couples to the elastic manifold and what type of LR model results. A general formulation of this coupling (see e.g. [3]) has the form:

$$V(x, u) = \int d^{D-d}z v(x, z) \rho(x, z, u), \quad (7)$$

where the defect potential lives in the D -dimensional space parameterized by (x, u) and $x \in R^d$ is the internal coordinate of the manifold. $\rho(x, z, u)$ is the manifold density. Each type of coupling to the disorder corresponds to a different function $\rho(x, z, u)$ and we now indicate the main cases.

1. Elastic interfaces in random-bond disorder

Let us first discuss elastic interfaces in the so-called Random Bond (RB) case, where the coupling between disorder and interface occurs only in the vicinity of the interface as e.g. for domain walls in magnets with random-bond disorder. This corresponds to the choice:

$$\rho(x, z, u) \sim \delta(z - u), \quad (8)$$

hence the additional variable z introduced in (7) is identical to u , the displacement field (with in general $D - d = N$). In that case:

$$V_{\text{RB}}(x, u) \sim v(x, u). \quad (9)$$

Consider now a uniform distribution of defects in the D -dimensional plane but *isotropically distributed within the (averaged direction) of the internal space of the manifold*. This

is given by (6) above with $D' = d$:

$$\overline{V_{\text{RB}}(x, u)V_{\text{RB}}(0, 0)} = g(x)R_2(u), \quad (10)$$

which is model (4) with a SR function $R_2(u)$ and, in full generality, $a = d - \varepsilon_d$. The physical realization in terms of extended defects is thus an interface ($d = 2$) in $D = 3$ with line defects all orthogonal to the u directions, isotropically distributed within the (average) plane of the interface, and $a = 1$. This is illustrated in Figure 1.

Another physical realization consists of extended defects with finite random lengths such that the distribution of lengths has a power-law tail for large lengths. For instance needles of variable lengths aligned along one direction could act on a directed polymer $d = 1$ as power-law correlated disorder in internal space.

An interesting, though qualitatively different case occurs when the extended defects are distributed isotropically in the whole (x, u) space. It will be discussed in Section IX. Finally note that we consider weak Gaussian disorder. It is possible that at strong disorder another phase exists where the line or manifold gets localized along the strongest extended defect.

2. Elastic interfaces in random field disorder

Random-Field (RF) disorder is described by the function:

$$\rho(x, z, u) \sim \Theta(u - z), \quad (11)$$

where $\Theta(z)$ is the Heaviside step function. This means that the change in energy when the interface moves between two configurations is proportional to the sum of all defect potentials in the volume (in R^D) spanned by this change. The discussion of the geometry of defects needed to produce LR disorder in internal space is identical to the last section. Substitution of Eq. (11) into Eq. (7) yields the RF disorder correlator which can be approximated by Eq. (4) with $R_i(u) \sim -u$ for large u .

3. Periodic systems

As an example of periodic systems we consider incommensurate single-Q CDWs. In that case $D = d$, hence the function $\rho(x, z, u) = \rho(x, u)$ in (7). The electron density of CDWs neglecting effects caused by an applied strain has the form [3, 6]:

$$\rho(x, \phi) \sim \rho_0 + \rho_1 \cos(2k_F(x_\perp - u(x))), \quad (12)$$

where the displacement $u(x)$ of the maximum of the density is related to the standard phase field via $\phi(x) = -2k_F u(x)$, where k_F is the Fermi wave-vector. The d -dimensional space is splitted into $x = (x_\parallel, x_\perp)$, with x_\perp denoting the modulation direction of the CDW and k_F the Fermi wave-vector.

We again consider the situation of extended defects all aligned with the direction x_\parallel and isotropically distributed in

that subspace. The random potential experienced by the CDW is given by

$$V(x, \phi) = h_1(x) \cos \phi(x) + h_2(x) \sin \phi(x), \quad (13)$$

with Gaussian distributed $h_1(x) = v(x) \cos(2k_F x_\perp)$ and $h_2(x) = v(x) \sin(2k_F x_\perp)$. On large scales $k_F x_\perp \gg 1$, and their cumulant can be approximated by (from (6)):

$$\overline{h_i(x)h_j(0)} = \frac{1}{2}v_{\text{SR}}^2\delta_{ij}\delta^d(x) + \frac{1}{2}\frac{v_{\text{LR}}^2}{x_\parallel^a}\delta_{ij}\delta(x_\perp), \quad (14)$$

where we have omitted all rapidly fluctuating contributions. Eqs. (13) and (14) give the potential correlator in a form which can be generalized to

$$\overline{V(x, u)V(x', u')} = R_1(u - u')\delta^d(x - x') + R_2(u - u')g(x_\parallel - x'_\parallel)\delta^{d_\perp}(x_\perp - x'_\perp), \quad (15)$$

with $d_\perp = 1$ and bare functions $R_i(\phi) = \frac{1}{2}v_i^2 \cos(\phi)$, $u \equiv \phi$. Thus periodic systems are described by periodic functions $R_i(u)$. Here d_\perp is the dimension of the transverse subspace. Note that the Hamiltonian $\mathcal{H}_{\text{XY}} = \int d^d x [\frac{1}{2}(\nabla\phi)^2 + V(x, \phi)]$ with $V(x, \phi)$ given by Eq. (13) and a Gaussian distribution of fields $\overline{h_i(x)h_j(x')} \sim g(x - x')$ describes the XY model with long-range correlated random fields. Therefore the latter can be mapped onto periodic manifolds with correlator (15) and $d_\perp = 0$, i.e. to model (4) with periodic functions $R_i(u)$. In the next section we will show how the FRG picture of model (15) can be obtained from the FRG results for model (4). It is worthy to note that in the case of periodic systems the integration in Fourier space is supposed to be over the first Brillouin zone. Note also that we have neglected the coupling of disorder to the long wavelength part of the density $-\rho_0 \int d^d x v(x) \nabla u(x)$ as it is usually irrelevant near the upper critical dimension. Indeed, in the replicated Hamiltonian (see below) this coupling generates additionally to the SR term $-1/T \int d^d x \sigma_1 \nabla u_a(x) \nabla u_b(x)$ the LR term

$$-\frac{1}{T} \int d^d x d^d x' \sigma_2 g(x_\parallel - x'_\parallel) \delta^{d_\perp}(x_\perp - x'_\perp) \nabla u_a(x) \nabla u_b(x').$$

For small disorder in the vicinity of the upper critical dimension both of them renormalize to zero according to

$$d_\ell \sigma_1 = (2 - d - 2\zeta)\sigma_1 + \dots, \quad (16)$$

$$d_\ell \sigma_2 = (2 - a - d_\perp - 2\zeta)\sigma_2 + \dots \quad (17)$$

IV. DYNAMICAL FORMALISM

The over-damped dynamics of the elastic manifold in a disordered medium can be described by the equation of motion

$$\eta \partial_t u_{xt} = c \nabla^2 u_{xt} + F(x, u_{xt}) + f_{xt}, \quad (18)$$

where η is the friction coefficient. In the presence of an applied force f the center of mass velocity is $v = L^{-d} \int_x \partial_t u_{xt}$.

The pinning force reads $F = -\partial_u V(x, u)$, and thus, for correlator (4) the second cumulant of the force is given by

$$\overline{F(x, u)F(x', u')} = \Delta_1(u - u')\delta^d(x - x') + \Delta_2(u - u')g(x - x'), \quad (19)$$

with $\Delta_i = -R_i''(u)$ in the bare model. In the following, we will always use $g(q) = |q|^{a-d}$, and $g(x) = \int_q e^{iqx} g(q)$.

The most important quantity of interest is the roughness exponent ζ measured in equilibrium or at the depinning transition $f = f_c$ defined by

$$C(x - x') = \overline{|u_x - u_{x'}|^2} \sim |x - x'|^{2\zeta}. \quad (20)$$

The velocity vanishes at the depinning transition as $v \sim |f - f_c|^\beta$, while the correlation length diverges at the transition as $\xi \sim |f - f_c|^{-\nu}$. One can also introduce the dynamic critical exponent z , which relates spatial and temporal correlations via $t \sim x^z$.

Let us briefly sketch how one can construct the perturbation theory in disorder. We adopt the dynamic formalism. It also allows us to obtain the statics equations (to one loop and $N = 1$ these can easily be deduced, as can be checked using replica). Instead of a direct solution of the equation of motion (18) with consequent averaging over different initial conditions and disorder configurations we employ the formalism of generating functional. Introducing the response field \hat{u}_{xt} we derive the effective action which reads

$$S = \int_{xt} i\hat{u}_{xt}(\eta\partial_t - c\nabla^2 + m^2)u_{xt} - \int_{xt} i\hat{u}_{xt}f_{xt}, \\ - \frac{1}{2} \int_{xtt'} i\hat{u}_{xt}i\hat{u}_{x't'}\Delta_1(u_{xt} - u_{x't'}) \\ - \frac{1}{2} \int_{xx'tt'} i\hat{u}_{xt}i\hat{u}_{x't'}g(x - x')\Delta_2(u_{xt} - u_{x't'}), \quad (21)$$

where we have added a small mass m , which plays the role of an IR cutoff. To study the critical domain one has to take the limit $m \rightarrow 0$. The average of the observable $A[u_{xt}]$ over dynamic trajectories with different initial conditions and over different disorder configurations can be written as follows

$$\langle A[u_{xt}] \rangle = \int \mathcal{D}[u]\mathcal{D}[\hat{u}] A[u_{xt}] e^{-S[u, \hat{u}]}. \quad (22)$$

Furthermore the response to the external perturbation f_{xt} , which is local in time and in space, can be computed using $\langle A[u_{xt}]i\hat{u}_{xt} \rangle = \frac{\delta}{\delta f_{xt}} \langle A[u_{xt}] \rangle$. Note that causality is fulfilled, and here we adopt the Ito convention, which results in getting rid of all closed loops composed of response functions.

In the absence of LR correlated disorder action, the Eq. (21) exhibits the so-called statistical tilt symmetry (STS), i.e. the invariance of the disorder terms under the tilt $u_{xt} \rightarrow u_{xt} + h_x$ with an arbitrary function h_x . The STS gives the exact identity $\int_t \mathcal{R}_{qt} = 1/cq^2$ for the response function $\mathcal{R}_{qt} = \langle u_{qt} i\hat{u}_{-q0} \rangle$, which implies that the elasticity is uncorrected by disorder to all orders. LR correlated disorder destroys the STS of action (21), and thus, in principle allows for a renormalization of the




FIG. 2: Diagrammatic rules: a - propagator, b - SR disorder vertex, c - LR disorder vertex; d and e one-loop diagrams generating the critical force at the depinning and giving correction to the mobility and elasticity.

elasticity. The quadratic part of the action (21) yields the free response function

$$\langle u_{qt} i\hat{u}_{-q0} \rangle_0 = R_{qt} = \frac{\Theta(t)}{\eta} e^{-(cq^2 + m^2)t/\eta}, \quad (23)$$

which can be used to generate the perturbation theory in disorder. The theory has two disorder interaction vertices $\Delta_1(u)$ and $\Delta_2(u)$. At each vertex $\Delta_i(u)$ there is one conservation rule for momentum and two for frequency while each vertex $\Delta_2(u)$ carries additional momentum dependence. In what follows we generalize the splitted diagrammatic method developed in Ref. [18], shown in Figure 2. As is the case for the model with SR-disorder, our model exhibits the so-called dimension reduction, both in the statics and in the dynamics. The naive perturbation theory obtained taking the functions $\Delta_i(u)$ analytic at $u = 0$ leads to the same result as that computed from the Gaussian theory setting $\Delta_i(u) = \Delta_i(0)$. In the limit $m \rightarrow 0$ the two-point function then read to all orders:

$$\overline{u_{qt}u_{-qt'}} = \frac{\Delta_1(0)}{c^2q^4} + \frac{\Delta_2(0)}{c^2q^{4+d-a}}. \quad (24)$$

The first term in Eq. (24) dominates in the limit $q \rightarrow 0$ for $a \geq d$, and LR disorder is irrelevant in this case, while the last term dominates for $a < d$. Eq. (24) results in $\zeta = (4 - d)/2$ for $a \geq d$ and $\zeta = (4 - a)/2$ for $a < d$ that are known to be incorrect. The physical reason for this is the existence of a large number of metastable states. The roughness exponent can be estimated using Flory arguments setting $u \sim x^\zeta$. Then the gradient term scales as $\nabla^2 u_x \sim x^{\zeta-2}$. The pinning force for SR disorder scales as $F(x, u_x) \sim x^{-(d+\zeta)/2}$ and for LR disorder as $F(x, u_x) \sim x^{-(a+\zeta)/2}$. Therefore in the regime where the behavior is governed by SR disorder the Flory estimate gives for RF disorder the Imry-Ma value $\zeta_{\text{SR}}^{\text{F}} = (4 - d)/3$ while for LR RF disorder we get $\zeta_{\text{LR}}^{\text{F}} = (4 - a)/3$. A similar argument constructed from the potential correlators $R_i(u)$ yields the Flory estimates $\zeta_{\text{SR}}^{\text{F}} = (4 - d)/5$ and $\zeta_{\text{LR}}^{\text{F}} = (4 - a)/5$ respectively, for the case of random-bond disorder. To obtain corrections to the Flory values, the FRG developed in Refs. [12, 14, 15, 16, 17, 18] will be employed. The solution is nontrivial because the renormalized disorder becomes non-analytic above the Larkin scale, and one has to deal with a non-analytic field theory. Here we generalize this approach to the case of LR correlated disorder.

V. FUNCTIONAL RENORMALIZATION

We now consider the renormalization of model (21). The subtleties arising for of the correlator (15) will shortly be discussed in the end. We carry out perturbation theory in the bare disorder correlators $\Delta_{i0}(u)$ and then introduce the renormalized correlators $\Delta_i(u)$. We will suppress the subscript "0" to avoid an overly complicated notation. According to the standard renormalization program we compute the effective action to one-loop order. Here we adopt the dimensional regularization of integrals and employ the minimal subtraction scheme to compute the renormalized quantities and absorb the poles in $\varepsilon = 4 - d$ and $\delta = 4 - a$ into multiplicative Z factors. When derivatives of the Δ_i at $u = 0$ occur, in the dynamics (i.e. at the depinning transition for dynamical quantities) they are taken at $u = 0^+$ as can be justified exactly for $N = 1$. In the statics the treatment is more subtle (as discussed in two loop studies [19]) but is not needed in the present one loop study.

Let us firstly consider the first order terms generated by expansion of e^{-S} in disorder. These terms are given by diagrams d and e shown in Figure 2. We start from

$$\int_{t>t',x} i\hat{u}_{xt}\Delta_1(u_{xt}-u_{x't'})i\hat{u}_{x't'} + \int_{t>t',x,x'} i\hat{u}_{xt}\Delta_2(u_{xt}-u_{x't'})g(x-x')i\hat{u}_{x't'}. \quad (25)$$

Expanding $\Delta_i(u)$ in a Taylor series and contracting one $i\hat{u}$ we obtain the leading corrections to the threshold force, friction and elasticity. The terms giving the threshold force to leading order are

$$\int_{t>t',x} i\hat{u}_{xt}\Delta_1'(0^+)R_{x=0,t-t'} + \int_{t>t',x,x'} i\hat{u}_{xt}\Delta_2'(0^+)g(x-x')R_{x-x',t-t'}. \quad (26)$$

They are strongly UV diverging ($\sim \Lambda^{d-2} + \Lambda^{a-2}$), and thus, are non-universal. The terms proportional to $\Delta_i''(0^+)$ can be rewritten as corrections to friction and elasticity using the expansion

$$u_{xt} - u_{x't'} = (t-t')\partial_t u_{xt} + (x-x')_i \frac{\partial}{\partial x_i} u_{xt} + (x-x')_i(x-x')_j \frac{1}{2} \frac{\partial^2 u_{xt}}{\partial x_i \partial x_j} + \mathcal{O}(\Delta t^2, \Delta x^3). \quad (27)$$

The first term in Eq. (27) gives the correction to friction

$$\begin{aligned} \delta\eta &= -\Delta_1''(0^+) \int_t t R_{x=0,t} - \Delta_2''(0^+) \int_{xt} t R_{x,t} g(x) \\ &= -\eta \left[\hat{\Delta}_1''(0^+) I_1 + \hat{\Delta}_2''(0^+) I_2 \right], \end{aligned} \quad (28)$$

where we have introduced $\hat{\Delta}_i(u) = \Delta_i(u)/c^2$. The one-loop integrals I_1 and I_2 diverge logarithmically and for $\varepsilon, \delta \rightarrow 0$




FIG. 3: 1-loop diagrams correcting disorder. The dot line corresponds to either SR disorder vertex (dash line) or to LR disorder vertex (wavy line). Diagrams of type a, b , and c contribute to SR disorder. Only diagrams of type d correct the LR disorder.

read

$$I_1 = \int_q \frac{1}{(q^2 + \hat{m}^2)^2} = K_d \frac{\hat{m}^{-\varepsilon}}{\varepsilon} + \mathcal{O}(1), \quad (29)$$

$$I_2 = \int_q \frac{q^{a-d}}{(q^2 + \hat{m}^2)^2} = K_d \frac{\hat{m}^{-\delta}}{\delta} + \mathcal{O}(1), \quad (30)$$

where we set $\hat{m} = m/\sqrt{c}$ and K_d is the area of a d -dimensional sphere divided by $(2\pi)^d$. To remove the poles in the mobility we introduce the corresponding Z -factor $\eta_R = Z_\eta^{-1}[\Delta_i]\eta$ which to one-loop order is given by

$$Z_\eta^{-1} = 1 - \hat{\Delta}_1''(0^+) I_1 - \hat{\Delta}_2''(0^+) I_2. \quad (31)$$

In the absence of LR correlated disorder the elasticity remains uncorrected to all orders due to the STS, while here the correction reads

$$\begin{aligned} \delta c &= \frac{1}{2d} \Delta_2''(0) \int_{xt} x^2 R_{x,t} g(x) \\ &= -\frac{1}{2d} \Delta_2''(0) \int_q g(q) \nabla_q^2 \frac{1}{cq^2 + m^2} \\ &= -c \frac{K_d}{d} \hat{\Delta}_2''(0) \frac{\varepsilon}{\delta} \hat{m}^{-\delta}. \end{aligned} \quad (32)$$

We have not set the second derivative at 0^+ as Δ_2 remains analytic as is discussed below. Furthermore the correction to elasticity (32) is finite for $\varepsilon, \delta \rightarrow 0$, and thus c does not acquire an anomalous dimension. However, we expect corrections at 2-loop order. If it is the case, one has to introduce a Z -factor which renormalizes elasticity: $c_R = Z_c^{-1}c$ with $Z_c = 1 + \mathcal{O}(\Delta_i^2)$.

In principal, due to the lack of STS, the KPZ term $\lambda(\nabla u_{xt})^2$ breaking the symmetry $u \rightarrow -u$ can be generated in the equation of motion (18) at the depinning transition. Indeed, diagram e in Figure 2, when expanding $\Delta(u)$ to second order in u , using (27), gives

$$\delta\lambda = \frac{1}{2d} \Delta_2'''(0^+) \int_{xt} x^2 R_{x,t} g(x). \quad (33)$$

Moreover the term with cubic symmetry ($M = 2$) and terms with higher order symmetries ($M > 2$) $\lambda_M \sum_i (\partial_i u_{xt})^{2M}$ can be generated by diagram e :

$$\delta\lambda_M = \frac{1}{d(2M)!} \Delta_2^{(2M+1)}(0^+) \int_{xt} R_{x,t} g(x) \sum_i x_i^{2M}. \quad (34)$$

However, as we will show later, if we start from bare analytic disorder distribution the LR disorder remains analytic along the FRG flow and the corresponding FP value $\Delta_2^*(u)$ is also an analytic function. Thus terms (33) and (34) are zero, provided that they are absent in the beginning. Moreover, the terms (34) are irrelevant in the RG sense for $M > 2$, [but not the KPZ term (33), see Ref. [46]]. This proves that our bare model (21) is a minimal model for the description of elastic manifolds in a random media with LR correlated disorder.

The corrections to disorder are given by the diagrams shown in Figure 3. The corresponding expressions read

$$\begin{aligned} \delta^1 \hat{\Delta}_1(u) = & -[\hat{\Delta}_1'(u)^2 + (\hat{\Delta}_1(u) - \hat{\Delta}_1(0))\hat{\Delta}_1''(u)]I_1 \\ & -[2\hat{\Delta}_1'(u)\hat{\Delta}_2'(u)^2 + (\hat{\Delta}_2(u) - \hat{\Delta}_2(0))\hat{\Delta}_1''(u) \\ & + \hat{\Delta}_1(u)\hat{\Delta}_2''(u)]I_2 - [\hat{\Delta}_2'(u)^2 + \hat{\Delta}_2(u)\hat{\Delta}_2''(u)]I_3, \end{aligned} \quad (35)$$

$$\delta^1 \hat{\Delta}_2(u) = -\hat{\Delta}_1(0)\hat{\Delta}_2''(u)I_1 - \hat{\Delta}_2(0)\hat{\Delta}_2''(u)I_2. \quad (36)$$

The one-loop integrals I_1 and I_2 have been defined in Eqs. (29) and (30), whereas I_3 is given by

$$I_3 = \int_q \frac{q^{2(a-d)}}{(q^2 + m^2)^2} = \frac{K_4 \hat{m}^{-2\delta+\varepsilon}}{2\delta - \varepsilon} + \mathcal{O}(1). \quad (37)$$

Let us define the renormalized dimensionless disorder Δ_i^R as

$$m^\varepsilon \Delta_1^R = \hat{\Delta}_1(u) + \delta^1 \hat{\Delta}_1(u), \quad (38)$$

$$m^\delta \Delta_2^R = \hat{\Delta}_2(u) + \delta^1 \hat{\Delta}_2(u). \quad (39)$$

The β_i functions are defined as the derivative of $\Delta_i^R(u)$ with respect to the mass m at fixed bare disorder $\Delta_i(u)$. In order to attain a fixed point, it is necessary to rescale the field u by m^ζ and write the β functions for the functions $\tilde{\Delta}_i := K_4 m^{-2\zeta} \Delta_i^R(um^\zeta)$:

$$\begin{aligned} \partial_\ell \tilde{\Delta}_1(u) = & (\varepsilon - 2\zeta)\tilde{\Delta}_1(u) + \zeta u \tilde{\Delta}_1'(u) \\ & - \frac{1}{2} \frac{d^2}{du^2} [\tilde{\Delta}_1(u) + \tilde{\Delta}_2(u)]^2 + A \tilde{\Delta}_1''(u), \end{aligned} \quad (40)$$

$$\partial_\ell \tilde{\Delta}_2(u) = (\delta - 2\zeta)\tilde{\Delta}_2(u) + \zeta u \tilde{\Delta}_2'(u) + A \tilde{\Delta}_2''(u), \quad (41)$$

where $A = [\tilde{\Delta}_1(0) + \tilde{\Delta}_2(0)]$, and $\partial_\ell := -m \frac{\partial}{\partial m}$.

The scaling behavior of the system is controlled by a stable fixed point $[\tilde{\Delta}_1^*(u), \tilde{\Delta}_2^*(u)]$ of flow equations (40) and (41). To determine the critical exponents let us start from power counting following Ref. [46]. The quadratic part of action (21) is invariant under $x \rightarrow xb$, $t \rightarrow tb^z$, $u \rightarrow ub^\zeta$, $\hat{u} \rightarrow \hat{u}b^{2-z-\zeta-d+\psi}$. Under this transformation the mobility, elasticity, and disorder scale at the Gaussian FP as $c \sim b^\psi$, $\eta \sim b^{2-z+\psi}$, $\tilde{\Delta}_1 \sim b^{4-d-2\zeta+2\psi}$ and $\tilde{\Delta}_2 \sim b^{4-a-2\zeta+2\psi}$. Thus SR disorder becomes relevant for $\zeta - \psi < (4-d)/2$ while LR disorder is naively relevant for $\zeta - \psi < (4-a)/2$.

Note that in the presence of STS $\psi = 0$ and we recover the conditions obtained at the end of Section III. The actual value of ζ will be fixed by the disorder correlators at the FP. The elasticity exponent ψ and the dynamic exponent z read

$$\psi = -m \frac{d}{dm} \ln Z_c(\tilde{\Delta}_i) \Big|_0, \quad (42)$$

$$z = 2 - \psi + m \frac{d}{dm} \ln Z_\eta(\tilde{\Delta}_i) \Big|_0, \quad (43)$$

where subscript "0" means a derivative at constant bare parameters. To one-loop order this yields

$$\psi = \mathcal{O}(\varepsilon^2, \varepsilon\delta, \delta^2), \quad (44)$$

$$z = 2 - \hat{\Delta}_1''(0) - \hat{\Delta}_2''(0). \quad (45)$$

The scaling relations then read [46]

$$\nu = \frac{1}{2 - \zeta + \psi}, \quad (46)$$

$$\beta = \nu(z - \zeta) = \frac{z - \zeta}{2 - \zeta + \psi}. \quad (47)$$

At zero velocity, the above calculation can be considered as a dynamical formulation of the equilibrium problem. However, one has to be careful with mapping the dynamic FRG equations to the static equations, because as shown in Ref. [19] the bare relation $\Delta_i = -R_i''(u)$ breaks down for the SR case at two-loop order. The standard derivation of the FRG equations in the statics is based on the renormalization of the replicated Hamiltonian. We have checked that similar to other systems with only SR disorder, the static FRG equations for systems with LR disorder can be obtained from the dynamic flow equations to one-loop order using the identification $\Delta_i = -R_i''(u)$. They read

$$\begin{aligned} \partial_\ell R_1(u) = & (\varepsilon - 4\zeta)R_1(u) + \zeta u R_1'(u) \\ & + \frac{1}{2} [R_1''(u) + R_2''(u)]^2 + A R_1''(u), \end{aligned} \quad (48)$$

$$\partial_\ell R_2(u) = (\delta - 4\zeta)R_2(u) + \zeta u R_2'(u) + A R_2''(u), \quad (49)$$

where $A = -[R_1''(0) + R_2''(0)]$.

In the case of the model with correlator given by Eq. (15), one has to distinguish between the transverse and parallel directions, and therefore introduce corresponding elastic coefficients c_\perp and c_\parallel . In the transverse direction, disorder is only δ -correlated and as a result the transverse elasticity is not corrected and can be set to 1. The power counting shows that the LR disorder is naively relevant for $\delta_1 = 4 - d_\perp - a < 0$. The one-loop integrals are logarithmically divergent and for $\varepsilon, \delta_1 \rightarrow 0$ are given by Eqs. (29), (30) and (37) with $\delta \rightarrow \delta_1$. Thus the above renormalization can be generalized to model (15) if one formally replaces $\delta \rightarrow \delta_1$.

Let us show how a non-analyticity of the disorder appears in the problem. We start from the bare analytic correlators with $\Delta_i''(0) < 0$. The flow equation for $y := -\Delta_1''(0) - \Delta_2''(0) \equiv R_1''''(0) + R_2''''(0) > 0$ reads

$$\partial_\ell y = \varepsilon y + 3y^2 + \gamma(m), \quad (50)$$

where $\gamma(m) = (\varepsilon - \delta)\Delta_2''(0)$. As we show below, the function $\hat{\Delta}_2(u)$ remains analytic along the whole FRG flow and at the fixed point (FP). The solution of Eq. (50) for any function $\gamma(m)$ bounded from below blows up at some finite scale m^* which can be associated with the inverse Larkin length. This blowup of y corresponds to the generation of a cusp singularity: $\Delta_1(u)$ becomes non-analytic at the origin and acquires for $m < m^*$ a non-zero $\Delta_1'(0^+)$. The precise estimation of the Larkin scale requires the solution of the pair of flow equations for both $\Delta_i(u)$.

Before studying different FPs, let us note an important property which is valid under all condition: if $\Delta_i(u)$ ($i = 1, 2$) is a solution of Eqs. (40) and (41) then $\kappa^2 \Delta_i(u/\kappa)$ is also a solution. Analogously, if $R_i(u)$ is a solution of Eqs. (48) and (49) then $\kappa^4 R_i(u/\kappa)$ is also a solution. We can use this property to fix the amplitude of the function in the non-periodic case, while for the periodic case the solution is unique as the period is fixed.

VI. NON-PERIODIC SYSTEMS: RANDOM BOND DISORDER

In this Section we study the scaling behavior of an elastic interface in a disordered environment with LR correlated RB disorder. To this aim we have to find a stationary solution (FP) of Eqs. (48) and (49) which decays exponentially fast at infinity as expected for RB disorder. The SR RB FP with $R_2(u) = 0$, which describes systems with only SR correlated disorder, was computed numerically in Refs. [12, 17, 18]. The corresponding roughness exponent to one-loop order is given by $\zeta_{\text{SR}} = 0.208298\varepsilon + \mathcal{O}(\varepsilon^2)$. We now look for a LR RB FP with $R_2(u) \neq 0$. Integrating Eq. (49) we obtain

$$\partial_\ell \int_0^\infty R_2(u) = (\delta - 5\zeta) \int_0^\infty R_2(u). \quad (51)$$

Therefore, the new LR RB FP, if it exists, has

$$\zeta_{\text{LRRB}} = \frac{\delta}{5}, \quad (52)$$

which will hold to all orders (since the RG equation for R_2 to higher orders can only be linear in $R_2(u)$ and involve even derivatives) and as a consequence, $\int_0^\infty du R_2(u)$ is exactly preserved along the FRG flow. Using our freedom to rescale $R_i(u)$, we introduce $\hat{\delta} := \delta/\varepsilon$, $R_i(u) =: \varepsilon r_i(u)$ and fix $r_1''(0) = -x$ and $r_2''(0) = -1$, where x is the parameter to be determined. The stationarity condition of Eqs. (48) and (49) reads

$$(1 - \frac{4}{5}\hat{\delta})r_1(u) + \frac{\hat{\delta}}{5}ur_1'(u) + \frac{1}{2}[r_1''(u) + r_2''(u)]^2 + (1+x)r_1''(u) = 0, \quad (53)$$

$$\frac{\hat{\delta}}{5}r_2(u) + \frac{\hat{\delta}}{5}ur_2'(u) + (1+x)r_2''(u) = 0. \quad (54)$$

TABLE I: Long-range correlated random bond fixed point. Shooting parameter $x = -r_1''(0)$, the maximal eigenvalue and the universal amplitude for different values of $\hat{\delta}$.

$\hat{\delta}$	$x(\hat{\delta})$	λ_1	$B(\hat{\delta})$
1.1	1.931986	—	33.89
1.2	1.121722	-0.160	31.37
1.3	0.922046	-0.262	31.64
1.4	0.825747	-0.365	32.41
1.5 ^a	0.766976	-0.469	33.34
2.0 ^b	0.639151	-1	38.44
3.0 ^c	0.562357	-2.120	48.10
∞	0.463619		∞

^aRandom lines in a planar interface ($d = 2$, $a = 1$).

^bRandom lines in a 3d manifold ($d = 3$, $a = 2$).

^cRandom planes in a 3d manifold ($d = 3$, $a = 1$).

Since Eq. (54) is linear in r_2 , it can be solved for fixed x by

$$r_2(u) = \frac{5(1+x)}{\hat{\delta}} \exp\left(-\frac{\hat{\delta}u^2}{10(1+x)}\right). \quad (55)$$

From the Taylor expansion of Eq. (53) around $u = 0$ we find

$$\begin{aligned} r_1(0) &= \frac{5(1-x^2)}{8\hat{\delta}-10}, \\ r_1'(0) &= 0, \end{aligned} \quad (56)$$

where the second condition excludes the possibility of a supercusp (the first line does not diverge since $x = 1$ for $\delta = 5/4$). Thus for fixed $\hat{\delta}$ the simultaneous equations (53) and (54) have a unique solution for any x but only for a specific x does the solution $r_1(u)$ decays exponentially fast to 0 for large u . To determine this value we employ the shooting method choosing x as our shooting parameter. For fixed x we integrate numerically Eq. (53) with $r_2(u)$ given by Eq. (55) from 0 to some large u_{max} with initial conditions (56). Then the shooting parameter x can be found by solving numerically the algebraic equation $r_1(u_{\text{max}}; x) = 0$. Increasing u_{max} we acquire the desired accuracy for x and $r_1(x)$. We were able to find the numerical solution with reasonable accuracy only for $\hat{\delta} \geq 1.1$. The typical FP functions $r_1^*(u)$ and $r_2^*(u)$ are shown in Figure 4. The actual values of x obtained by shooting for different $\hat{\delta}$ are summarized in Table I.

Let us now check the stability of SR and LR FPs. To that end we linearize the FRG equations about each FP. In the vicinity of a FP, the linearized flow equations have solutions which are pure power-laws in m , i.e. scale as $m^{-\lambda}$ with a discrete spectrum of eigenvalues λ . A stable fixed point has all eigenvalues $\lambda < 0$. Substituting $r_i \rightarrow r_i^*(u) + z_i(u)$ into the flow equations and keeping only terms which are linear in $z_i(u)$, we derive the linearized flow equations at the FP




FIG. 4: Fixed point describing the interface in a medium with long-range correlated random bond disorder (LR RB FP) for $\hat{\delta} = 2$. The SR part $r_1^*(u)$ is a non-analytic function with $r_1^{*''}(0^+) \neq 0$. The LR part $r_2^*(u)$ is an analytic function. Here we report minus their second derivative.

$\{r_1^*(u), r_2^*(u)\}$:

$$\begin{aligned} (1 - 4\zeta_1)z_1(u) + \zeta_1 u z_1'(u) + [r_1^{*''}(u) + r_2^{*''}(u)] \\ \times [z_1''(u) + z_2''(u)] + (1+x)z_1''(u) \\ + A_0 r_1^{*''}(u) = \lambda z_1(u), \end{aligned} \quad (57)$$

$$\begin{aligned} (\hat{\delta} - 4\zeta_1)z_2(u) + \zeta_1 u z_2'(u) \\ + (1+x)z_2''(u) + A_0 r_2^{*''}(u) = \lambda z_2(u), \end{aligned} \quad (58)$$

where we have introduced $\zeta = \varepsilon\zeta_1$, $A_0 = -[z_1''(0) + z_2''(0)]$, and λ is also measured in unites of ε . Note that because of the freedom to rescale $r_i(u)$ we always have the eigenmode $z_i^{(0)}$ with marginal eigenvalue $\lambda_0 = 0$. As shown in Ref. [45] for SR RB FP the corresponding eigenfunction is given by $z_1^{(0)} = u r_1^{*'}(u) - 4r_1^*(u)$, $z_2^{(0)} = 0$, while the next eigenvalue $\lambda_1 = -1$ corresponds to $z_1^{(1)} = \zeta_1^{\text{SR}} u r_1^{*'}(u) + (1 - 4\zeta_1^{\text{SR}})r_1^*(u)$, $z_2^{(1)} = 0$. Here $\{r_1^*, r_2^* = 0\}$ is the SR RB FP and the Taylor expansion of the function r_1^* can be found in Ref. [45]. Thus the SR RB FP is stable in the SR disorder subspace ($r_2 = z_2 = 0$). Let us check its stability with respect to introduction of LR correlated disorder. From Eq. (58) follows that the maximal eigenvalue $\lambda_{\text{max}} = \hat{\delta} - 5\zeta_1^{\text{SR}}$ corresponds to the exponential eigenfunction $z_2(u) = \exp(-\zeta_1^{\text{SR}} u^2 / |2r_1^{*''}(0)|)$ with $r_1^{*''}(0) = -0.577$ for SR RB FP. As a consequence, the LR correlated disorder destabilizes the SR RB FP if $\hat{\delta} > 5\zeta_1^{\text{SR}} \approx 1.041$, or equivalently, using (52) if $\zeta^{\text{SR}} < \zeta^{\text{LR}}$. This criterion was of course expected.

We now check the stability of the LR RB FP $\{r_1^*(u), r_2^*(u) \neq 0\}$. It also has a marginal eigenvalue $\lambda_0 = 0$ with eigenfunctions given by $z_i^{(0)} = u r_i^{*'}(u) - 4r_i^*(u)$ that can be checked by direct substitution into Eq. (57) and (58). Eq. (58) allows for an analytical solution which reads

$$z_2(u) = \frac{5A_0}{2\hat{\delta} + 5\lambda} \left[\frac{\hat{\delta} u^2}{5(1+x)} - 1 \right] \exp\left(-\frac{\hat{\delta} u^2}{10(1+x)}\right) \quad (59)$$




FIG. 5: Second derivative of eigenfunctions $z_1(u)$ and $z_2(u)$ computed at the LR RB FP for $\hat{\delta} = 2$.

We are free to fix the length of the eigenvectors, for instance, by the condition $z_2''(0) = 1$ which gives

$$A_0 = \frac{1}{3\hat{\delta}}(1+x)[2\hat{\delta} + 5\lambda]. \quad (60)$$

Thus to find the eigenvalue λ and the eigenfunction z_1 we have to solve Eq. (57) with condition $z_1''(0) = -1 - A_0$ and require an exponentially fast decay of the solution at large u . The only case for which we succeeded to construct the solution analytically is $\hat{\delta} = 2$, which is depicted in Figure 5. It has eigenvalue $\lambda = -1$ and reads

$$z_1(u) = -\frac{1}{3}u r_1^{*'}(u) + \frac{1}{2}r_1^*(u) + \frac{5}{6}r_2^*(u), \quad (61)$$

$$z_2(u) = -\frac{1}{3}[u r_2^{*'}(u) + r_2^*(u)]. \quad (62)$$

For other values of $\hat{\delta}$ we solve Eq. (57) numerically using λ as a shooting parameter and require an exponentially fast decay of $z_1(u)$ for large u . To compute the numerical solution we need the initial conditions. Expanding Eq. (57) in a Taylor series, we obtain

$$z_1(0) = \frac{5[x^2(2\hat{\delta} + 5\lambda) + 5x(\hat{\delta} + \lambda) + 3\hat{\delta}]}{3\hat{\delta}(5 - 4\hat{\delta} - 5\lambda)}, \quad (63)$$

$$z_1'(0) = 0. \quad (64)$$

Apart from the marginal eigenvalue $\lambda_0 = 0$, the largest eigenvalue is λ_1 . It is shown for different $\hat{\delta} > 1.1$ in Table I. The negative sign of λ_1 reflects the stability of the LR RB FP. For $\hat{\delta} \leq 1.1$ we failed to compute the numerical solution with reasonable accuracy. However, the largest eigenvalue computed at LR RB FP λ_1 tends to 0 for $\hat{\delta} \rightarrow 1.1$ and the SR RB FP becomes unstable for $\hat{\delta} > 1.041$ with respect to LR correlated disorder. Thus we expect that the LR RB FP is stable for $\hat{\delta} > 1.041$. Moreover, the largest eigenvalue within accessible accuracy is well approximated by $\lambda_1 = 0.1917(\zeta_1^{\text{SR}} - \zeta_1^{\text{LR}})$ that gives $\lambda_1 = -0.06$ for $\hat{\delta} = 1.1$.

Besides the roughness exponent there is another universal quantity which is of interest. This is the displacement correlation function, which behaves like

$$\overline{u_q u_{-q}} = \mathcal{A}_d q^{-(d+2\zeta)}. \quad (65)$$

Let us show that in contrast to systems with only SR-correlated disorder, this system, whose behavior is controlled by the LR RB FP, has a universal amplitude \mathcal{A}_d . Indeed, according to equation (51), the integral $\int du R_2^{\text{tr}}(u)$ is preserved along the flow and is fixed to its bare value Q , where we have introduced the actual renormalized correlator R_2^{tr} which is connected to R_2 given by Eq. (55) by the relation $R_i^{\text{tr}} = \kappa^4 R_i(u/\kappa)$ with κ given by

$$\kappa = \frac{Q^{1/5}}{(2\pi)^{1/10}} \left(\frac{\hat{\delta}}{5(1+x)} \right)^{3/10}, \quad (66)$$

where we used $\int du R_2^{\text{tr}}(u) = Q$. Then the amplitude can be written to one-loop order as follows [19]

$$\begin{aligned} \mathcal{A}_d &= \frac{1}{K_4} (-R_1^{\text{tr}''}(0) - R_2^{\text{tr}''}(0)) \\ &= \frac{\kappa^2}{K_4} (1+x) = Q^{2/5} B(\hat{\delta}), \end{aligned} \quad (67)$$

where we have introduced the universal function

$$B(\hat{\delta}) = \frac{8\pi^2}{(2\pi)^{1/5}} (1+x(\hat{\delta}))^{2/5} \left(\frac{\hat{\delta}}{5} \right)^{3/5}. \quad (68)$$

Values for $x(\hat{\delta})$ and for $B(\hat{\delta})$ for different $\hat{\delta}$ are shown in Table I.

VII. NON-PERIODIC SYSTEMS: RANDOM FIELD DISORDER

We now address the problem of an elastic interface in a medium with LR-correlated RF disorder. We expect that similar to systems with uncorrelated disorder this universality class also describes the depinning transition. To see that systems with RB disorder flow in the dynamics to the RF FP, one has to include either effects of a finite velocity or consider two-loop contributions, that go beyond of the scope of the present work; but we expect the mechanism to be the same as in Ref. [18].

Let us look for a solution of Eqs. (40) and (41), which decays exponentially fast at infinity as expected for RF disorder. From Eq. (41) it follows that

$$\partial_\ell \int_0^\infty \Delta_2(u) = (\delta - 3\zeta) \int_0^\infty \Delta_2(u). \quad (69)$$

Therefore $\int_0^\infty \Delta_2(u)$ is preserved along the FRG flow fixing the roughness exponent to

$$\zeta_{\text{LRRF}} = \frac{\delta}{3} + O(\varepsilon^2, \delta^2, \varepsilon\delta), \quad (70)$$

which coincides with the Flory estimate. Introducing $\Delta_i(u) = \varepsilon y_i(u)$, $\zeta = \varepsilon\zeta_1$ and fixing $y_1(0) = x$, $y_2(0) = 1$ we can rewrite the stationary form of Eqs. (40) and (41) as follows ($\zeta_1 = \hat{\delta}/3$):

$$\begin{aligned} (1 - 2\zeta_1)y_1(u) + \zeta_1 u y_1'(u) - \frac{1}{2} \frac{d^2}{du^2} [y_1(u) + y_2(u)]^2 \\ + [1+x]y_1''(u) = 0, \end{aligned} \quad (71)$$

$$(\hat{\delta} - 2\zeta_1)y_2(u) + \zeta_1 u y_2'(u) + [1+x]y_2''(u) = 0. \quad (72)$$

Eq. (72) can be solved analytically giving

$$y_2(u) = \exp\left(-\frac{\hat{\delta}u^2}{6(1+x)}\right). \quad (73)$$

Substituting the FP function (73) in Eq. (71), we obtain a closed differential equation for the function $y_1(u)$. Expanding around $u = 0$ we find

$$y_1'(0) = -\frac{1}{3} \sqrt{9x + 3\hat{\delta} - 6x\hat{\delta}}, \quad (74)$$

$$y_1''(0) = \frac{1}{3} - \frac{\hat{\delta}x - 2}{9x + 1}, \quad (75)$$

$$y_2'(0) = 0, \quad (76)$$

$$y_2''(0) = -\frac{\hat{\delta}}{3(x+1)}. \quad (77)$$

Thus we can compute numerically the solution $y_1(u)$ for any fixed $\hat{\delta}$ and $y_1(0) \equiv x$, however, only for special values of x this solution decays exponentially at infinity. The corresponding solution can be computed using the shooting method as described above, using x as a shooting parameter (see Table II).

A pair of typical FP functions is shown in Figure 6. Surprisingly, the function $y_1(u)$ obtained by shooting satisfies $\int_0^\infty du y_1(u) = 0$, characteristic for RB type correlations along the u direction. In other words, the LR RF FP is in fact of mixed type: RB for the SR part and RF for the LR part of the disorder correlator. This can be understood as follows: Consider the flow of $\int_0^\infty du y_1(u)$. It is obtained by integrating the l.h.s. of Eq. (71) from 0 to infinity, and by inserting $\zeta_1 = \hat{\delta}/3$:

$$\begin{aligned} \partial_\ell \int_0^\infty du y_1(u) &= (1 - \hat{\delta}) \int_0^\infty du y_1(u) \\ &+ \frac{1}{2} \frac{d}{du} [y_1(u) + y_2(u)]^2 \Big|_{u=0} - (1+x)y_1'(0), \end{aligned} \quad (78)$$

where we have used the fact that most terms in the FRG-equation (71) are total derivatives. Finally, we remark that the second line of Eq. (78) cancels exactly provided that $y_2(u)$ is an analytical function, leaving us with

$$\partial_\ell \int_0^\infty du y_1(u) = (1 - \hat{\delta}) \int_0^\infty du y_1(u). \quad (79)$$

This means that for LR-correlated disorder, i.e. $\hat{\delta} > 1$, the integral of y_1 indeed scales to 0. A non-trivial fixed point is

TABLE II: Long-range correlated random field disorder. Shooting parameter $x = y_1(0)$, the maximal eigenvalue and the universal amplitude for different values of $\hat{\delta}$.

$\hat{\delta}$	$x(\hat{\delta})$	λ_1	$B(\hat{\delta})$
1.1	0.562872	-0.1	140.43
1.2	0.525082	-0.2	142.23
1.3	0.496948	-0.3	144.27
1.4	0.475110	-0.4	146.44
1.5 ^a	0.457638	-0.5	148.66
2.0 ^b	0.404989	-1.0	159.66
3.0 ^c	0.362329	-2.0	179.04

^aRandom lines in planar interface ($d = 2, a = 1$).

^bRandom lines in a 3d manifold ($d = 3, a = 2$).

^cRandom planes in a 3d manifold ($d = 3, a = 1$).

possible at 2-loop order for depinning. We remind that in [18] it has been shown that at 2-loop order and for SR-correlated disorder, new terms arise in the FRG-equation, which do *not* integrate to 0. Indeed, this is the mechanism which leads to a break-down of the result $\zeta_{\text{SRRF}} = \varepsilon/3$, at depinning. The same terms will appear here. We expect that the additional diagrams due to LR-correlations do not exactly cancel these terms, especially since these terms are proportional to the derivative at the cusp, and LR-disorder will probably remain analytique, thus not contribute to the anomalous terms. These considerations let us expect that at 2-loop order the integral of $y_1(u)$ will be small, but non-zero.

Let us finally check the stability of the SR RF FP and new LR RF FP. At the SR RF FP the roughness is given by $\zeta_{\text{SRRF}} = \varepsilon/3$, and thus, we expect the crossover from the SR universality to LR at $\delta > \varepsilon$, which follows from the condition $\zeta_{\text{SRRF}} = \zeta_{\text{LRRF}}$. To check the stability of the FPs, we follow the strategy used for the RB case and linearize the flow equations about the RF FPs. We obtain

$$(1 - 2\zeta_1)z_1(u) + \zeta_1 u z_1'(u) - \frac{d^2}{du^2} \left\{ [y_1^*(u) + y_2^*(u)] \times [z_1(u) + z_2(u)] \right\} + (1+x)z_1''(u) + A_0 y_1^{*''}(u) = \lambda z_1(u), \quad (80)$$

$$(\hat{\delta} - 2\zeta_1)z_2(u) + \zeta_1 u z_2'(u) + (1+x)z_2''(u) + A_0 y_2^{*''}(u) = \lambda z_2(u), \quad (81)$$

where we have introduced $A_0 = z_1(0) + z_2(0)$. Firstly we prove our conclusion on the stability of SR RF FP with respect to LR correlated disorder. To that end we solve Eq. (58) assuming that $\zeta_1 = \zeta_1^{\text{SR}} = 1/3$ and $x = y_1^{\text{SR}}(0) = 2/9$. We obtain that $z_2 = \exp(-\zeta_1^{\text{SR}} u^2 / (2y_1^{\text{SR}}))$ and the corresponding eigenvalue $\lambda_{\text{max}} = \hat{\delta} - 3\zeta_1^{\text{SR}}$. Therefore, indeed the SR RF FP becomes unstable with respect to LR disorder for $\delta > \varepsilon$.

We now focus on the stability of the LR RF FP. Analysis of the linearized flow equations (80) and (81) shows that there is at least one eigenvector $z_i^{(0)} = u\Delta_i^*(u) - 2\Delta_i^*$ with marginal eigenvalue $\lambda_0 = 0$, which corresponds to the free-




FIG. 6: Fixed point describing the interface in a medium with long-range correlated random field disorder (LR RF FP) for $\hat{\delta} = 2$. The SR correlator $y_1^*(u)$ has a cusp at origin and formally corresponds to RB type of correlation in direction u . The LR correlator $y_2^*(u)$ is an analytic function of RF type.

dom of rescaling. For arbitrary λ , Eq. (81) can be solved analytically:

$$z_2(u) = \frac{A_0 (\delta u^2 - 3(1+x))}{3(2\delta + 3\lambda)(1+x)^2} \exp\left(-\frac{\hat{\delta} u^2}{6(1+x)}\right). \quad (82)$$

We are free to fix the length of the eigenvectors, for instance, by the condition $z_2(0) = 1$ which gives

$$A_0 = -\frac{1}{\hat{\delta}}(2\hat{\delta} + 3\lambda)(1+x). \quad (83)$$

Thus to find the eigenvalue λ and the eigenfunction z_1 we have to solve Eq. (80) with condition $z_1(0) = A_0 - 1$ and require an exponentially fast decay for large u . We need the initial conditions which can be found by expanding Eq. (80) in a Taylor series:

$$z_1(0) = -1 - \frac{1}{\hat{\delta}}(2\hat{\delta} + 3\lambda)(1+x), \quad (84)$$

$$z_1'(0) = -\left\{ [9\lambda^2 + \lambda(12\hat{\delta} - 9)](x+1) + \hat{\delta}^2(4x+7) - \hat{\delta}(6x+9) \right\} / 2\hat{\delta}\sqrt{9x-6x\hat{\delta}+3\hat{\delta}}. \quad (85)$$

$\hat{\delta} = 2$ is the only case in which we succeeded to find a completely analytical solution. It has $\lambda_1 = -1$ and reads

$$z_1(u) = u y_1^{*'}(u) - \frac{1}{2} y_1^*(u) - \frac{3}{2} y_2^*(u), \quad (86)$$

$$z_2(u) = u y_2^{*'}(u) + y_2^*(u). \quad (87)$$

For other values of $\hat{\delta}$, Eq. (87) remains correct, while to obtain $z_1(u)$, we solve Eq. (80) numerically, using λ as a shooting parameter. As can be seen from Table II, the largest eigenvalue satisfies $\lambda_1(\hat{\delta}) = 1 - \hat{\delta} \equiv 3(\zeta_1^{\text{SR}} - \zeta_1^{\text{LR}})$. This result can

be obtained analytically as follows. Integrating Eq. (80) from 0 to ∞ , we get the condition

$$\int_0^\infty du z_1(u) (1 - \hat{\delta} - \lambda) = 0, \quad (88)$$

proving that as long as $\int z_1(u) \neq 0$, one has $\lambda = 1 - \hat{\delta}$. Therefore the LR RF FP is stable for $\delta > \varepsilon$. Inserting this value of λ back into (80), we obtain after some simplifications

$$\begin{aligned} 0 &= \frac{\hat{\delta}}{3} \frac{d}{du} (u z_1(u)) - \frac{d^2}{du^2} [Y(u) z_1(u) + W(u)], \\ Y(u) &:= y_1(u) - y_1(0) + y_2(u) - y_2(0), \\ W(u) &:= z_2(u) [y_1(u) + y_2(u)] - y_1(0) [z_1(0) + z_2(0)]. \end{aligned} \quad (89)$$

The first equation can be integrated with the result

$$\frac{\hat{\delta}}{3} u z_1(u) = \frac{d}{du} [Y(u) z_1(u) + W(u)]. \quad (90)$$

This is equivalent to

$$z_1 \left[Y' - \frac{\hat{\delta}}{3} u \right] + z_1' Y = -W'. \quad (91)$$

The homogenous equation reads

$$[\ln z_1 Y]' = \frac{\hat{\delta}}{3} \frac{u}{Y}. \quad (92)$$

Its solution is

$$z_1(u) = \frac{C}{Y(u)} \exp \left[\frac{\hat{\delta}}{3} \int_0^u ds \frac{s}{Y(s)} \right] \quad (93)$$

with some constant C . A solution to the inhomogeneous equation is obtained by replacing C by $C(u)$, and inserting the latter into Eq. (91). This yields

$$C(u) = - \int_0^u dt W'(t) \exp \left[-\frac{\hat{\delta}}{3} \int^t ds \frac{s}{Y(s)} \right]. \quad (94)$$

Putting together everything, we obtain

$$z_1(u) = - \frac{1}{Y(u)} \int_0^u dt W'(t) \exp \left[-\frac{\hat{\delta}}{3} \int_u^t ds \frac{s}{Y(s)} \right]. \quad (95)$$

Let us now compute the universal amplitude defined by Eq. (65). According to Eq. (69) the integral $\int du \Delta_2(u)$ is preserved along the FRG flow and can be fixed by its bare value Q . The relation between the actual renormalized disorder Δ_2^{tr} and the rescaled disorder $\Delta_2(u)$ given by Eq. (73) reads $\Delta_i^{\text{tr}} = \kappa^2 \Delta_i(u/\kappa)$. We obtain

$$\kappa = \frac{Q^{1/3}}{\varepsilon^{1/3}} \left(\frac{2\hat{\delta}}{3\pi(1+x)} \right)^{1/6}, \quad (96)$$




FIG. 7: Eigenfunctions $z_1(u)$ and $z_2(u)$ computed at the LR RF FP for $\hat{\delta} = 2$.

where we have fixed $\int du \Delta_2^{\text{tr}}(u) = Q$. Then the amplitude can be written as

$$\begin{aligned} \mathcal{A}_d &= \frac{1}{K_4} (\Delta_1^{\text{tr}}(0) + \Delta_2^{\text{tr}}(0)) \\ &= \frac{\varepsilon \kappa^2}{K_4} (1+x) = \varepsilon^{1/3} Q^{2/3} B(\hat{\delta}), \end{aligned} \quad (97)$$

with the universal functions given by

$$B(\hat{\delta}) = 8\pi^2 (1+x)^{2/3} \left(\frac{2\hat{\delta}}{3\pi} \right)^{1/3}. \quad (98)$$

Values of $B(\hat{\delta})$ for different $\hat{\delta}$ are shown in Table II.

Depinning. We are now in the position to study the depinning transition, which we expect is controlled by the LR RF FP. The dynamic critical exponent z defined by Eq. (45) is given to one-loop order by

$$z = 2 - \frac{\varepsilon}{3} + \frac{\delta}{9} + \mathcal{O}(\varepsilon^2, \delta^2, \varepsilon\delta), \quad (99)$$

where we have used Eqs. (75) and (77) which give $y_1''(0) + y_2''(0) = 1/3 - \hat{\delta}/9$. Other exponents can be computed using scaling relations (47) and (46), for example

$$\beta = 1 - \frac{\varepsilon}{6} + \frac{\delta}{18} + \mathcal{O}(\varepsilon^2, \delta^2, \varepsilon\delta). \quad (100)$$

It is remarkable that for $\delta > 3\varepsilon$ the exponent β is larger than 1, and z larger than 2. This seems to imply some different physics - yet to be understood - in the avalanche process which makes the motion slower near depinning than in the SR case. The analyticity of Δ_2 seems to suggest some smoother motion at large scale, while short scale motion remains jerky and avalanche like. Finally, note that at the SR RF FP $z_{\text{SR}} = 2 - 2\varepsilon/9$, $\beta_{\text{SR}} = 1 - \varepsilon/9$, and thus, the exponents are continuous functions of ε and δ .

VIII. PERIODIC SYSTEMS

We now study periodic systems with disorder correlator given by Eq. (4), which we can refer to as an XY model with LR correlated defects. The results for CDWs with LR correlated disorder defined by correlator (15) can then be obtained by substituting $\delta \rightarrow \delta_1 = 4 - d_\perp - a$. It is sufficient to consider the system with the period fixed to 1, since other systems can be related to the latter using the freedom to rescale. As a consequence the roughness exponent for periodic systems is $\zeta = 0$. At variance with interfaces we introduce reduced parameters according to $\Delta_i(u) = \delta y_i(u)$, $A = y_1(0) + y_2(0)$ and $\hat{\varepsilon} = \varepsilon/\delta$. Then the fixed point equations can be written as follows

$$\hat{\varepsilon} y_1(u) - \frac{1}{2} \frac{d^2}{du^2} [y_1(u) + y_2(u)]^2 + A y_1''(u) = 0, \quad (101)$$

$$y_2(u) + A y_2''(u) = 0. \quad (102)$$

Equation (102) can be solved analytically. Its solution is

$$y_2 = y_2(0) \cos(2\pi u), \quad A = 1/(2\pi)^2. \quad (103)$$

Equation (101) can be solved analytically for $\hat{\varepsilon} = 0$:

$$y_1 = y_1(0) + y_2(0) (1 - \cos(2\pi u)) - \frac{1}{2\pi} \sqrt{2y_2(0) (1 - \cos(2\pi u))}. \quad (104)$$

The coefficients $y_i(0)$ are determined by potentiality of the Δ_i , i.e. from the conditions

$$\int_0^1 du y_1(u) = \int_0^1 du y_2(u) = 0, \quad (105)$$

and the identity $y_1(0) + y_2(0) = 1/(2\pi)^2$. They read

$$y_1(0) = 1/(2\pi)^2 - 1/64, \quad (106)$$

$$y_2(0) = 1/64. \quad (107)$$

For $\hat{\varepsilon} > 0$ Eq. (101) can be written in the following form

$$\hat{\varepsilon} y_1(u) - \frac{1}{2} \frac{d^2}{du^2} \left\{ [y_1(u) + y_2(u)]^2 - \frac{y_1(u)}{\pi^2} \right\} = 0, \quad (108)$$

where $y_2(u)$ is given by Eq. (103) with $y_2(0) = 1/(2\pi)^2 - y_1(0)$. Expanding Eq. (108) in a Taylor series about $u = 0$ we find that $y_1'(0) = -\sqrt{1/(2\pi)^2 - y_1(0)(1 - \hat{\varepsilon})}$. Thus for any fixed $0 \leq \hat{\varepsilon} < 1$ and $y_1(0)$ we have only one solution $y_1(u)$, but only for a specific $y_1(0)$ this solution fulfills the condition $y_1(1) = y_1(0)$. To find this value we employ the shooting method using $y_1(0)$ as a shooting parameter. The values of $y_1(0)$ computed for different $\hat{\varepsilon}$ are summarized in Table III. The corresponding eigenfunctions $y_1(u)$ are depicted in Figure 8.

While the roughness exponent is zero, the system forms a Bragg glass phase with a slow growth of the displacements according to

$$\overline{(u_x - u_0)^2} = \mathcal{A}_d \ln |x|, \quad (109)$$

TABLE III: Periodic systems with LR correlated disorder. The shooting parameter: $y_1(0)$ and two first eigenvalues for different $\hat{\varepsilon}$.

$\hat{\varepsilon}$	$y_1(0)$	λ_1	λ_2
0	0.00971	0.0	—
1/3	0.01089	0.333	-4.089
1/2	0.01183	0.500	-0.500
2/3 ^a	0.01348	0.667	-0.280
0.8	0.01645	0.800	-0.125
0.9	0.02346	0.900	-0.013

^aCorresponds to $d = 2$, $a = 1$, i.e. line defects (e.g. dislocations) along the plane of a CDW.




FIG. 8: Fixed point of a periodic system with LR correlated disorder. The SR disorder correlator $y_1(u)$ computed for different values of $\hat{\varepsilon}$.

where \mathcal{A}_d is a universal amplitude, which to one-loop order is given by

$$\mathcal{A}_d^{(\text{LR})} = \frac{2K_d}{K_4} [\Delta_1^*(0) + \Delta_2^*(0)] = \frac{\delta}{2\pi^2}, \quad (110)$$

where we have restored the factor of $1/K_4$ previously absorbed in Δ_i . The SR periodic FP is characterized by $\mathcal{A}_d^{(\text{SR})} = \varepsilon/18 + \mathcal{O}(\varepsilon^2)$. It is interesting to compare (110) with the prediction of the Gaussian variational approximation for the SR disorder case $\mathcal{A}_{d,\text{GVA}}^{(\text{SR})} = \varepsilon/(2\pi^2)$. We expect the crossover between LR and SR FPs at $\mathcal{A}_d^{(\text{LR})} = \mathcal{A}_d^{(\text{SR})}$, i.e. LR disorder to be relevant for

$$\frac{\varepsilon}{\delta} < \frac{9}{\pi^2} \approx 0.912. \quad (111)$$

We now check the stability of the SR and LR periodic FPs. The flow equations linearized about the FP read

$$\begin{aligned} \hat{\varepsilon} z_1(u) - \frac{d^2}{du^2} \{ [y_1^*(u) + y_2^*(u)] [z_1(u) + z_2(u)] \} \\ + A z_1''(u) + A_0 y_1''(u) = \lambda z_1(u), \end{aligned} \quad (112)$$

$$z_2(u) + A z_2''(u) + A_0 y_2''(u) = \lambda z_2(u), \quad (113)$$

where we have introduced $A_0 = z_1(0) + z_2(0)$. Let us remind that the SR FP is unstable with respect to non-potential perturbations even in the subspace of SR disorder. Indeed, the SR periodic FP

$$\Delta_1^*(u) = \frac{\varepsilon}{6} \left[\frac{1}{6} - u(1-u) \right], \quad \Delta_2^*(u) = 0 \quad (114)$$

has in the SR subspace the positive eigenvalue $\lambda_1 = 1$, corresponding to the non-potential eigenfunction $z_1 = 1$. All other eigenfunctions are potential, i.e. fulfill condition (105), and have negative eigenvalues [45]. If we add LR correlated disorder, the solution of Eq. (113) yields

$$z_2(u) = \cos 2\pi u. \quad (115)$$

The corresponding eigenvalue $\lambda_{\text{SR}} = 1 - \hat{\varepsilon}\pi^2/9$ confirms our estimation for the stability of the SR periodic FP (111). For the LR periodic FP we still have Eq. (115) with

$$A_0 = -\frac{\lambda}{1 - 4\pi^2 y_1(0)}, \quad z_1(0) = A_0 - 1. \quad (116)$$

Equation (112) has a periodic solution only for a discrete set of eigenvalues λ_i (the first two are shown in Table III). It follows from the Table that $\lambda_1 = \hat{\varepsilon} > 0$. In analogy with the SR periodic FP the LR periodic FP is unstable with respect to a non-potential perturbation corresponding to λ_1 . The latter is obtained by integrating (113) over one period,

$$(\hat{\varepsilon} - \lambda) \int_0^1 du z_1(u) = 0. \quad (117)$$

As long as the integral does not vanish, this gives the reported eigenvalue $\lambda_1 = \hat{\varepsilon}$. Indeed as it can be seen from Figure 9 we have $\int_0^1 du z_1^{(1)}(u) \neq 0$ and $\int_0^1 du z_1^{(n)}(u) \neq 0$ for $n \geq 2$.

Depinning. We now focus on the depinning transition of the periodic system with LR correlated disorder. At the LR periodic FP we have

$$y_1''(0) = 1 + \frac{\hat{\varepsilon}}{3} - 4\pi^2 y_1(0), \quad (118)$$

$$y_2''(0) = -4\pi^2 y_2(0), \quad (119)$$

and thus, $y_1''(0) + y_2''(0) = \hat{\varepsilon}/3$. The dynamic critical exponent z defined by Eq. (45) reads to one-loop order

$$z^{\text{LR}} = 2 - \frac{\varepsilon}{3} + \mathcal{O}(\varepsilon^2, \delta^2, \varepsilon\delta), \quad (120)$$

Therefore for periodic systems $z^{\text{LR}} = z^{\text{SR}}$ to one-loop order.

IX. FULLY ISOTROPIC EXTENDED DEFECTS

In this Section we briefly examine the effect of a defect distribution isotropic in the whole (x, u) space. Consider first interfaces in random-bond type disorder. From (5) and (9) one finds:

$$R(x, u) = \overline{V_{\text{RB}}(x, u) V_{\text{RB}}(0, 0)} \sim \frac{v_{\text{LR}}^2}{|x^2 + u^2|^{a/2}}, \quad (121)$$




FIG. 9: Two first eigenvectors computed at the LR periodic FP. Eigenfunctions $z_1^{(1)}(u)$ and $z_1^{(2)}(u)$ for $\hat{\varepsilon} = 2/3$.

and thus, the u and x dependencies are coupled in the bare correlator. For the present discussion we consider N , the number of components of u , arbitrary, hence $D = N + d$. We recall that $a = D - \varepsilon_d$. The question to which universality class this model belongs is subtle. It turns out that it does not correspond to LR disorder in internal space, but rather SR disorder in internal space and LR disorder in the u direction, hence $R_2 = 0$ but $R_1(u)$ long range in u . To see this let us consider at fixed u the integral $\int d^d x R(x, u)$. We can distinguish two cases:

(i): for $a > d$, this integral is convergent at large x , hence we clearly have SR disorder in the x -direction, and $R_1(u) \sim |u|^{d-a}$ at large u . This however is LR disorder in u . This case has been studied using FRG and yields, for $a < a_c(d, N)$ a roughness exponent exactly given by the Flory value $\zeta(a, d) = (4 - d)/(4 + a - d)$. The value $a_c(d, N)$ can be estimated using the value for the SR disorder roughness exponent, by requiring $\zeta(a_c(d, N), d) = \zeta_{\text{SR, RB}}(d, N)$ (small deviations can arise as discussed in [7]).

(ii): for $a < d$, the situation is more subtle and one may be tempted to argue, since $\int d^d x R(x, u)$ diverges in the infrared, that disorder LR in x is produced. This is however not the case, as can be seen on the Fourier transform $R(q, P)$, where P is the momentum associated to u , and q to x . One has $R(q, P) \sim (q^2 + P^2)^{(a-d-N)/2}$, which has a well defined limit $R(q = 0, P) = P^{a-d-N}$. This corresponds again, as we argue, to a SR correlator in space with $R_1(u) - R_1(0) \sim |u|^{d-a}$. As is often the case LR models require some trivial subtractions. The subtracted correlator $R(x, u) - R(x, 0)$ has indeed a convergent integral $\sim |u|^{d-a}$ at large u , while subtracting a u -independent piece does not change the model. The critical case $a = d$ is described by the logarithmic model $R(x, u) - R(x, 0) \sim \ln |u|$ which has $\zeta = (4 - d)/4$ in all dimensions [47].

To summarize, isotropic distributions of defects isotropic in the (x, u) space also yield LR models, but not of the type (4) studied here. For isotropic line defects one finds $\zeta = (4 - d)/(3 + N)$ (i.e. $\zeta = 3/4$ for a directed polymer in $D = 1 + 1$,




FIG. 10: 2d domain wall moving in 3D magnet with fully isotropic planar defects.

$\zeta = 3/5$ in $D = 1 + 2$, and for an interface $D = 2 + 1$, $\zeta = 2/5$). Isotropic planar defects yield $\zeta = (4-d)/(2+N)$, hence $\zeta = 2/3$ for a $(D = 2 + 1)$ -dimensional interface. This case is illustrated in Figure 10. Note that in that case there are infinitely many lines of defects inside the interface with random directions (the intersections of the planar defects with the interface gives lines), but that this does not suffice to create power-law correlations in internal space, as can be seen from the example where the planar defects are orthogonal to the interface.

Finally in the periodic case, such as for CDWs, isotropic disorder in the full space $(x_{\parallel}, x_{\perp})$ again leads to correlations (15), but now the function $g(x_{\parallel} - x'_{\parallel})$ decays exponentially beyond a length scale set by the disorder period (as can be seen in Fourier space considering the discrete P modes). Hence the problem is described by the standard (SR) random periodic class.

X. CONCLUSION

We have studied elastic interfaces and periodic systems in a medium with LR correlated disorder, both in equilibrium and

at the depinning transition. This type of long-range correlations exists in the internal space of the manifold, and we have discussed how it can be realized in terms of extended defects, or anisotropic defects with a broad distribution of lengths. Using a dynamic formalism we derived the FRG flow equations for the SR and LR parts of the disorder correlator and found three new FPs, which describe three new universality classes. All new FPs are characterized by a non-analytic SR part of the disorder correlator and an analytic LR part. We have computed the corresponding exponents and universal amplitudes in a double expansion in $\epsilon = 4 - d$ and $\delta = 4 - a$. For RB type of disorder we find that the LR correlation of disorder is relevant for $\delta > 1.041\epsilon$ and results in the roughness exponent $\zeta = \delta/5$, while for $\delta < 1.041\epsilon$ the scaling behavior is controlled by the SR RB FP with $\zeta = 0.208298\epsilon$. We find that the presence of RF disorder results in a mixed FP with the SR correlator corresponding formally to RB type of disorder and an analytic RF LR correlator. The LR RF FP which is also expected to control the depinning transition is stable for $\delta > \epsilon$ giving $\zeta = \delta/3$ and $\beta = 1 - \epsilon/6 + \delta/18$. The LR correlated periodic FP is stable for $\epsilon < 0.912\delta$ and gives a slow logarithmic growth of displacements with universal amplitude $\mathcal{A}_d^{\text{LR}} = \delta^2/(2\pi^2)$. It is remarkable that this type of disorder yields an exponent β for the velocity-force characteristics which can be larger than unity and a dynamical exponent larger than 2. This striking behavior might be relevant for experiments, and gives a strong motivation for numerical studies of the problem, e.g. to understand the nature of motion at the depinning transition in these systems.

Acknowledgments

We thank Dima Feldman for useful discussions. AAF acknowledges support from the European Commission through a Marie Curie Postdoctoral Fellowship under contract number MIF1-CT-2005-021897. PLD and KJW are supported by Agence Nationale de la Recherche under contract number 05-BLAN-0099-01.

-
- [1] D. S. Fisher, Phys. Rep. **301**, 113 (1998).
 - [2] M. Kardar, Phys. Rep. **301**, 85 (1998).
 - [3] S. Brazovskii and T. Nattermann, Adv. Phys. **53**, 177 (2004).
 - [4] S. Lemerle, J. Ferre, C. Chappert, V. Mathet, T. Giamarchi, and P. Le Doussal, Phys. Rev. Lett. **80**, 849 (1998).
 - [5] D. Wilkinsion and J.F. Willemsen, J. Phys. A **16**, 3365 (1983).
 - [6] G. Grüner, Rev. Mod. Phys. **60**, 1129 (1988).
 - [7] G. Blatter, M.V. Feigel'man, V.B. Geshkenbein, A.I. Larkin, and V.M. Vinokur, Rev. Mod. Phys. **66**, 1125 (1994).
 - [8] T. Nattermann and S. Scheidl, Adv. Phys. **49**, 607 (2000).
 - [9] T. Giamarchi and P. Le Doussal, Phys. Rev. Lett. **72**, 1530 (1994); Phys. Rev. B **52**, 1242 (1995).
 - [10] M. Mezard and G. Parisi J. Phys. A **23**, L1229 (1990).
 - [11] S.E. Korshunov, Phys. Rev. B **48**, 3969 (1993).
 - [12] D. S. Fisher, Phys. Rev. Lett. **56**, 1964 (1986).
 - [13] A.I. Larkin, Sov. Phys. JETP **31**, 784 (1970).
 - [14] T. Nattermann, S. Stepanow, L.-H. Tang, and H. Leschhorn, J. Phys. II France **2**, 1483 (1992).
 - [15] H. Leschhorn, T. Nattermann, S. Stepanow, and L.-H. Tang, Ann. Phys. (Leipzig) **6**, 1 (1997).
 - [16] O. Narayan and D. S. Fisher, Phys. Rev. B **48**, 7030 (1993).
 - [17] P. Chauve, P. Le Doussal, and K.J. Wiese, Phys. Rev. Lett. **86**, 1785 (2001).
 - [18] P. Le Doussal, K.J. Wiese, and P. Chauve, Phys. Rev. B **66**, 174201 (2002).
 - [19] P. Le Doussal, K.J. Wiese, and P. Chauve, Phys. Rev. E **69**, 026112 (2004).
 - [20] P. Le Doussal and K.J. Wiese, Phys. Rev. B **68**, 174202 (2003).

- [21] L. Roters and K.D. Usadel, Phys. Rev. E **65**, 027101 (2002)
- [22] L. Roters, S. Lübeck, and K. D. Usadel, Phys. Rev. E **66**, 026127 (2002).
- [23] A. Rosso, A.K. Hartmann, and W. Krauth, Phys. Rev. E **67**, 021602 (2003).
- [24] A.A. Middleton, P. Le Doussal, and K.J. Wiese, cond-mat/0606160 (preprint); P. Le Doussal, cond-mat/0605490 (preprint).
- [25] P. Chauve, T. Giamarchi, and P. Le Doussal, Phys. Rev. B **62**, 6241 (2000).
- [26] S.N. Dorogovtsev, Phys. Lett. **76A**, 169 (1980).
- [27] D. Boyanovsky and J.L. Cardy, Phys. Rev. B **26**, 154 (1982).
- [28] I.D. Lawrie and V.V. Prudnikov, J. Phys. C **17**, 1655 (1984).
- [29] A.A. Fedorenko, Phys. Rev. B **69**, 134301 (2004).
- [30] A. Weinrib and B.I. Halperin, Phys. Rev. B **27**, 413 (1983).
- [31] E. Korutcheva and F. Javier de la Rubia, Phys. Rev. B **58**, 5153 (1998).
- [32] V.V. Prudnikov, P.V. Prudnikov, and A.A. Fedorenko, J. Phys. A **32**, L399 (1999); Phys. Rev. B **62**, 8777 (2000).
- [33] L. Civale, A.D. Marwick, T.K. Worthington, M.A. Kirk, J.R. Thompson, L. Krusin-Elbaum, Y. Sun, J.R. Clem, and F. Holtzberg, Phys. Rev. Lett. **67**, 648 (1991).
- [34] D.R. Nelson and V.M. Vinokur, Phys. Rev. Lett. **68**, 2398 (1992).
- [35] T. Emig and T. Nattermann, cond-mat/0604345 (preprint).
- [36] T. Hwa, P. Le Doussal, D.R. Nelson, and V.M. Vinokur, Phys. Rev. Lett. **71**, 3545 (1993).
- [37] L. Civale, L. Krusin-Elbaum, J.R. Thompson, R. Wheeler, A.D. Marwick, M.A. Kirk, Y.R. Sun, F. Holtzberg, and C. Feild, Phys. Rev. B **50**, 4102 (1994).
- [38] T. Hwa, D.R. Nelson, and V.M. Vinokur, Phys. Rev. B **48**, 1167 (1993).
- [39] P. Chauve, Ph.D. thesis, Université de Paris XI Orsay, 2000; <http://www.lpt.ens.fr/~chauve/>
- [40] P. Chauve, P. Le Doussal, and T. Giamarchi Phys. Rev. B **61**, R11906(2000).
- [41] Y. Yamazaki, A. Holz, M. Ochiai, and Y. Fukuda, Physica A **150**, 576 (1988).
- [42] C. Vasquez R., R. Paredes V., A. Hasmy, and R. Jullien, Phys. Rev. Lett. **90**, 170602 (2003).
- [43] D.E. Feldman and R.A. Pelcovits, Phys. Rev. E **70**, 040702(R) (2004).
- [44] E. Medina, T. Hwa, M. Kardar, and Y.-C. Zhang, Phys. Rev. A **39**, 3053 (1989).
- [45] P. Le Doussal and K.J. Wiese, Phys. Rev. E **68**, 046118 (2003).
- [46] P. Le Doussal and K.J. Wiese, Phys. Rev. E **67**, 016121 (2003).
- [47] P. Le Doussal, Phys. Rev. Lett. **96**, 235702 (2006) and to be published.